

NEUMANN ISOPERIMETRIC CONSTANT ESTIMATE FOR CONVEX DOMAINS

XIANZHE DAI, GUOFANG WEI, AND ZHENLEI ZHANG

ABSTRACT. We present a geometric and elementary proof of the local Neumann isoperimetric inequality on convex domains of a Riemannian manifold with Ricci curvature bounded below.

1. INTRODUCTION

Isoperimetric and Sobolev inequalities are equivalent inequalities (see e.g. Theorem 1.3 below) which play important role in geometric analysis on manifolds. Indeed, doing analysis on manifolds usually depends on the estimate of the Sobolev constant which could then be obtained via the isoperimetric constant. There are extensive work on isoperimetric constant estimates. An important method pioneered by Gromov relies on the geometric measure theory and its regularity theory, which works for closed manifolds or convex domains with smooth boundary, see e.g. the survey article [9] and the recent paper [14]. One may also obtain an estimate through Li-Yau gradient estimate for heat kernel [12] and the equivalence of heat kernel bounds, Sobolev inequality, isoperimetric inequality, see [15, Page 448], which again requires smooth and convex boundary. Another method using needle decomposition from convex geometry has also been very successful and, very recently, been combined with optimal transport and extended to non-smooth case, see [3] and the references therein. For star-shaped domains in a manifold with Ricci curvature bounded from below, Buser [2] gave an elementary proof for a Neumann isoperimetric constant (the Cheeger constant) estimate using comparison geometry, but the estimate depends on the in and out radius of the domain, which does not give a uniform estimate for convex domain as the in-radius might be small. We would like to point out that for convex domains with non-smooth boundaries, the estimate for isoperimetric constant is only obtained in the very recent paper [3] mentioned above. In this short note we give a very geometric and elementary proof of a Neumann isoperimetric inequality, albeit with non-optimal constant, for convex domains whose boundaries need not be smooth.

First we recall some definitions.

1991 *Mathematics Subject Classification.* 53C20.

XD is partially supported by the Simons Foundation, NSF and NSFC, GW partially supported by the Simons Foundation and NSF DMS 1506393, ZZ partially supported by CNSF11371256 .

Definition 1.1. When M is compact (with or without boundary), the Neumann α -isoperimetric constant of M is defined by

$$\text{IN}_\alpha(M) = \sup_H \frac{\min\{\text{vol}(M_1), \text{vol}(M_2)\}^{1-\frac{1}{\alpha}}}{\text{vol}(H)},$$

where H varies over compact $(n-1)$ -dim submanifold of M which divides M into two disjoint open submanifolds M_1, M_2 (with or without boundary).

Definition 1.2. The Neumann α -Sobolev constant of M is defined by

$$\text{SN}_\alpha(M) = \sup_{f \in C^\infty(M)} \frac{\inf_{a \in \mathbb{R}} \|f - a\|_{\frac{\alpha}{\alpha-1}}}{\|\nabla f\|_1}.$$

The isoperimetric constant and Sobolev constant are equivalent.

Theorem 1.3 ([4], see also [11]). *For all $n \leq \alpha \leq \infty$,*

$$\text{IN}_\alpha(M) \geq \text{SN}_\alpha(M) \geq \frac{1}{2} \text{IN}_\alpha(M).$$

For convenience we consider the normalized Neumann α -isoperimetric and α -Sobolev constant:

$$\text{IN}_\alpha^*(M) = \text{IN}_\alpha(M) \text{vol}(M)^{1/\alpha}, \quad \text{SN}_\alpha^*(M) = \text{SN}_\alpha(M) \text{vol}(M)^{1/\alpha}.$$

Using comparison geometry and Vitali covering we give an estimate on the normalized Neumann isoperimetric constant for convex domain in terms of the Ricci curvature lower bound and the diameter of the domain.

Theorem 1.4. *Let (M, g) be a complete Riemannian manifold of dimension n , with $\text{Ric} \geq -(n-1)K$ for some $K \geq 0$. Let Ω be a bounded convex domain. Then*

$$(1.1) \quad \text{IN}_n^*(\Omega) \leq 40^n e^{11(n-1)\sqrt{K}d} \cdot d$$

where d is the diameter of the domain Ω . In particular, if M is closed with diameter d , then

$$(1.2) \quad \text{IN}_n^*(M) \leq 40^n e^{11(n-1)\sqrt{K}d} \cdot d.$$

Corollary 1.5. *Let (M, g) be a complete Riemannian manifold of dimension n , with nonnegative Ricci curvature. Let Ω be a bounded convex domain. Then*

$$(1.3) \quad \text{IN}_n^*(\Omega) \leq 40^n \cdot d$$

where d is the diameter of the domain Ω . In particular, if M is closed with diameter d , then

$$(1.4) \quad \text{IN}_n^*(M) \leq 40^n \cdot d.$$

Remark 1.6. The case when Ω equals the whole manifold is well-known. The reference we mentioned earlier for convex domain in the literature deals with domains with (smooth) convex boundary which is a stronger condition.

Remark 1.7. For balls we can obtain both Dirichlet and Neumann isoperimetric constant estimates even under the much weaker integral Ricci lower bound assumption [7, 18]. On the other hand it is not clear if that will remain true for convex domains.

Remark 1.8. Using the mean curvature estimate from [16] one gets similar estimate when the Bakry-Emery Ricci curvature is bounded from below and oscillation of the potential function is bounded.

2. PROOF OF THEOREM 1.4

The proof goes by a covering argument of Anderson [1], combined with an observation of Gromov [10]. See [1] or [7] for a similar argument of estimating the local Dirichlet isoperimetric constant. First of all we recall a lemma whose proof is a slight modification of Gromov's observation [10, 5.(C)].

Lemma 2.1. *Let M^n be a complete Riemannian manifold. Let Ω be a convex domain of M and H be any hypersurface dividing Ω into two parts Ω_1, Ω_2 . For any Borel subsets $W_i \subset \Omega_i$, there exists x_1 in one of W_i , say W_1 , and a subset W in another one, W_2 , such that*

$$(2.1) \quad \text{vol}(W) \geq \frac{1}{2} \text{vol}(W_2)$$

and any $x_2 \in W$ has a unique minimal geodesic connecting to x_1 which intersects H at some z such that

$$(2.2) \quad \text{dist}(x_1, z) \geq \text{dist}(x_2, z).$$

The convexity assumption of Ω is essential. It implies that any minimal geodesic with endpoints in different parts must intersects H . The Bishop-Gromov relative volume comparison theorem gives

Lemma 2.2. *Let H , W and x_1 be as in the lemma above. Then*

$$(2.3) \quad \text{vol}(W) \leq 2^{n-1} D e^{(n-1)\sqrt{K}D} \text{vol}(H')$$

where $D = \sup_{x \in W} \text{dist}(x_1, x)$ and H' is the set of intersection points with H of geodesics $\gamma_{x_1, x}$ for all $x \in W$.

Proof. Let $\Gamma \subset S_{x_1}$ be the set of unit vectors such that $\gamma_v = \gamma_{x_1, x_2}$ for some $x_2 \in W$. We compute the volume in the polar coordinate at x_1 . Write $dv = \mathcal{A}(\theta, t) d\theta \wedge dt$ in the polar coordinate $(\theta, t) \in S_{x_1} \times \mathbb{R}^+$. For any $\theta \in \Gamma$, let $r(\theta)$ be the radius such that $\exp_{x_1}(r\theta) \in H$. Then $W \subset \{\exp_{x_1}(r\theta) | \theta \in \Gamma, r(\theta) \leq r \leq 2r(\theta)\}$. So, by

relative volume comparison,

$$\begin{aligned}
\text{vol}(W) &\leq \int_{\Gamma} \int_{r(\theta)}^{2r(\theta)} \mathcal{A}(\theta, t) dt d\theta \\
&\leq \frac{\sinh^{n-1}(2\sqrt{K}D)}{\sinh^{n-1}(\sqrt{K}D)} \int_{\Gamma} r(\theta) \mathcal{A}(\theta, r(\theta)) d\theta \\
&\leq D \frac{\sinh^{n-1}(2\sqrt{K}D)}{\sinh^{n-1}(\sqrt{K}D)} \text{vol}(H').
\end{aligned}$$

The required estimate follows from $\frac{\sinh(2t)}{\sinh t} = 2 \cosh t \leq e^t$ whenever $t \geq 0$. \square

Corollary 2.3. *Let H be any hypersurface dividing a convex domain Ω into two parts Ω_1, Ω_2 . For any ball $B = B_r(x)$ we have*

$$(2.4) \quad \min(\text{vol}(B \cap \Omega_1), \text{vol}(B \cap \Omega_2)) \leq 2^{n+1} r e^{(n-1)\sqrt{K}d} \text{vol}(H \cap B_{2r}(x))$$

where $d = \text{diam}(\Omega)$. In particular, if $B \cap \Omega$ is divided equally by H , we have

$$(2.5) \quad \text{vol}(B_r(x) \cap \Omega) \leq 2^{n+2} r e^{(n-1)\sqrt{K}d} \text{vol}(H \cap B_{2r}(x))$$

Proof. Put $W_i = B \cap \Omega_i$ in the above lemma and notice that $D \leq 2r$ and $H' \subset H \cap B_{2r}(x)$. \square

Now we are ready to prove our main theorem.

Proof of Theorem 1.4. We may assume that $\text{vol}(\Omega_1) \leq \text{vol}(\Omega_2)$. For any $x \in \Omega_1$, let r_x be the smallest radius such that

$$\text{vol}(B_{r_x}(x) \cap \Omega_1) = \text{vol}(B_{r_x}(x) \cap \Omega_2) = \frac{1}{2} \text{vol}(B_{r_x}(x) \cap \Omega).$$

Let $d = \text{diam}(\Omega)$. By above corollary,

$$(2.6) \quad \text{vol}(B_{r_x}(x) \cap \Omega) \leq 2^{n+2} r_x e^{(n-1)\sqrt{K}d} \text{vol}(H \cap B_{2r_x}(x)).$$

The domain Ω_1 has a covering

$$\Omega_1 \subset \bigcup_{x \in \Omega_1} B_{2r_x}(x).$$

By Vitali Covering Lemma, cf. [13, Section 1.3], we can choose a countable family of disjoint balls $B_i = B_{2r_{x_i}}(x_i)$ such that $\bigcup_i B_{10r_{x_i}}(x_i) \supset \Omega_1$. Applying the relative

volume comparison theorem and the convexity of Ω we have

$$\begin{aligned}
\text{vol}(\Omega_1) &\leq \sum_i \frac{\int_0^{10r_{x_i}} \sinh^{n-1}(\sqrt{K}t) dt}{\int_0^{r_{x_i}} \sinh^{n-1}(\sqrt{K}t) dt} \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega_1) \\
&\leq 10 \sum_i \frac{\sinh^{n-1}(10\sqrt{K}r_{x_i})}{\sinh^{n-1}(\sqrt{K}r_{x_i})} \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega_1) \\
&\leq 10 \frac{\sinh^{n-1}(10\sqrt{K}d)}{\sinh^{n-1}(\sqrt{K}d)} \sum_i \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega_1) \\
&\leq 10^n e^{9(n-1)\sqrt{K}d} \sum_i \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega_1) \\
&= 2^{-1} \cdot 10^n \cdot e^{9(n-1)\sqrt{K}d} \sum_i \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega).
\end{aligned}$$

Moreover, since the balls B_i are disjoint, (2.6) gives,

$$\text{vol}(H) \geq \sum_i \text{vol}(B_i \cap H) \geq 2^{-n-2} e^{-(n-1)\sqrt{K}d} \sum_i r_{x_i}^{-1} \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega).$$

These two estimates lead to

$$\begin{aligned}
\frac{\text{vol}(\Omega_1)^{\frac{n-1}{n}}}{\text{vol}(H)} &\leq 2 \cdot 20^n e^{10(n-1)\sqrt{K}d} \frac{(\sum_i \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega))^{\frac{n-1}{n}}}{\sum_i r_{x_i}^{-1} \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega)} \\
&\leq 40^n e^{10(n-1)\sqrt{K}d} \frac{\sum_i \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega)^{\frac{n-1}{n}}}{\sum_i r_{x_i}^{-1} \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega)} \\
&\leq 40^n e^{10(n-1)\sqrt{K}d} \sup_i \frac{\text{vol}(B_{r_{x_i}}(x_i) \cap \Omega)^{\frac{n-1}{n}}}{r_{x_i}^{-1} \text{vol}(B_{r_{x_i}}(x_i) \cap \Omega)} \\
&= 40^n e^{10(n-1)\sqrt{K}d} \sup_i \left(\frac{r_{x_i}^n}{\text{vol}(B_{r_{x_i}}(x_i) \cap \Omega)} \right)^{\frac{1}{n}}.
\end{aligned}$$

On the other hand, since $\text{vol}(\Omega_1) \leq \text{vol}(\Omega_2)$, we have $r_x \leq d$ for any $x \in \Omega_1$. Thus, by the relative volume comparison and convexity of Ω again, we have

$$\text{vol}(\Omega) \leq \frac{\int_0^d \sinh^{n-1}(\sqrt{K}t) dt}{\int_0^{r_x} \sinh^{n-1}(\sqrt{K}t) dt} \text{vol}(B_{r_x}(x) \cap \Omega).$$

Therefore,

$$\text{vol}(\Omega)^{\frac{1}{n}} \cdot \frac{\text{vol}(\Omega_1)^{\frac{n-1}{n}}}{\text{vol}(H)} \leq 40^n e^{10(n-1)\sqrt{K}d} \sup_{0 < r \leq d} \left(\frac{r^n \int_0^d \sinh^{n-1}(\sqrt{K}t) dt}{\int_0^r \sinh^{n-1}(\sqrt{K}t) dt} \right)^{\frac{1}{n}}.$$

The last term on the right hand side has the estimate

$$\frac{r^n \int_0^d \sinh^{n-1}(\sqrt{K}t) dt}{\int_0^r \sinh^{n-1}(\sqrt{K}t) dt} \leq r^n \cdot \frac{d}{r} \cdot \frac{\sinh^{n-1}(\sqrt{K}d)}{\sinh^{n-1}(\sqrt{K}r)} \leq d^n \cdot \frac{\sinh^{n-1}(\sqrt{K}d)}{(\sqrt{K}d)^{n-1}} \leq d^n e^{(n-1)\sqrt{K}d}.$$

The required normalized Neumann isoperimetric constant estimate now follows. \square

REFERENCES

- [1] M. T. Anderson, *The L^2 structure of moduli spaces of Einstein metrics on 4-manifolds*, GAFA., 2 (1992), 29-89.
- [2] P. Buser, *A note on the isoperimetric constant*, Ann. Sci. École Norm. Sup. (4) 15 (1982), no. 2, 213 - 230.
- [3] F. Cavalletti & A. Mondino *Sharp and rigid isoperimetric inequalities in metric-measure spaces with lower Ricci curvature bounds*, arXiv:1502.06465 [math.MG], to appear in *Inventiones*.
- [4] J. Cheeger, *A lower bound for the smallest eigenvalue of the Laplacian*, in *Problems in Analysis*, R. Gunning ed., 195-199. Princeton University Press, 1970.
- [5] J. Cheeger and S.T. Yau, *A lower bound for the heat kernel*, Comm. Pure Appl. Math., 34 (1981), 465-480.
- [6] S. Y. Cheng & S. T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*, Comm. Pure Appl. Math. 28 (1975), 333-354.
- [7] X. Dai, G. Wei and Z. Zhang, *Local Sobolev Constant Estimate for Integral Ricci Curvature Bounds*, arXiv:1601.08191.
- [8] S. Gallot, *Isoperimetric inequalities based on integral norms of Ricci curvature*, Soc. Math. de France, Asterisque 157-158 (1988), 191-216.
- [9] S. Gallot, *Inegalites isoperimetriques et analytiques sur les varietes riemanniennes*. (French) [Isoperimetric and analytic inequalities on Riemannian manifolds] On the geometry of differentiable manifolds (Rome, 1986). Asterisque No. 163-164 (1988), 31-91.
- [10] M. Gromov, *Paul Levy's isoperimetric inequality*, Publications IHES, 1980.
- [11] P. Li, *Geometric Analysis*, Cambridge studies in advanced mathematics 134, Cambridge University Press, 2012.
- [12] P. Li and S.T. Yau, *On the parabolic kernel of the Schrodinger operator*, Acta Math. 156 (1986) 153- 201.
- [13] F.H. Lin and X.P. Yang, *Geometric Measure Theory-An Introduction*, Science Press, Beijing/New York and Intern. Press, Boston, 2002.
- [14] E. Milman, *Sharp isoperimetric inequalities and model spaces for the curvature-dimension-diameter condition*, J. Eur. Math. Soc. (JEMS) 17 (2015), no. 5, 1041 - 1078.
- [15] L. Saloff-Coste, *Uniformly elliptic operators on Riemannian manifolds* Journal of Diff. Geom. 36 (1992), 417-450.
- [16] G. Wei and W. Wylie, *Comparison Geometry for the Bakry-Emery Ricci Tensor*, Journal of Diff. Geom. 83, no. 2 (2009), 377-405.
- [17] D. Yang, *Convergence of Riemannian manifolds with integral bounds on curvature. I*, Ann. Scient. Éc. Norm. Sup., 25 (1992), 77-105.
- [18] Z. Zhang, *Notes on the isoperimetric constant estimate*, preprint.

(Xianzhe Dai) DEPARTMENT OF MATHEMATICS, ECNU, SHANGHAI AND UCSB, SANTA BARBARA CA 93106, EMAIL: DAI@MATH.UCSB.EDU

(Guofang Wei) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA CA 93106, EMAIL: WEI@MATH.UCSB.EDU

(Zhenlei Zhang) DEPARTMENT OF MATHEMATICS, CAPITAL NORMAL UNIVERSITY, CHINA, EMAIL: ZHLEIGO@ALIYUN.COM