

Bases

The Punch Line: We have an efficient way to define subspaces using collections of vectors in them.

Warm-Up: Are these sets linearly independent? What do they span?

(a) $\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \right\} \subset \mathbb{R}^3$

(b) All vectors in \mathbb{R}^{42} with a zero in at least one component

(c) $\left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 2 \\ -3 \\ 2 \end{bmatrix} \right\} \subset \mathbb{R}^4$

(d) $\{1, t-1, (t-1)^2 + 2(t-1)\} \subset \mathcal{P}_2$

Bases: A *basis* for a vector space is a linearly independent spanning set. Every finite spanning set contains a basis by removing linearly dependent vectors, and many finite linearly independent sets may be extended to be a basis by adding vectors (if eventually this process terminates in a spanning set).

1 Are these sets bases for the indicated vector spaces? If not, can vectors be removed (which?) or added (how many?) to make it a basis?

(a) $\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \right\} \subset \mathbb{R}^3$

(c) $\left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 2 \\ -3 \\ 2 \end{bmatrix} \right\} \subset \mathbb{R}^4$

(b) $\left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \end{bmatrix} \right\} \subset \mathbb{R}^2$

(d) $\{(t-1), (t-1)^2, (t-1)^3\} \subset \mathcal{P}_3$

Finding Bases in \mathbb{R}^n : We're often interested in subspaces of the form $\text{Nul } A$ and $\text{Col } A$ for some matrix A . Fortunately, we can extract both by examining the Reduced Echelon Form of A .

A basis for $\text{Col } A$ consists of all columns in A itself which correspond to pivot columns in the REF of A . A basis for $\text{Nul } A$ consists of the vector parts corresponding to each free variable in a parametric vector representation of the solution set of the homogeneous equation $A\vec{x} = \vec{0}$, which we can find from the REF of A . Caution: In general, although free variables correspond to non-pivot columns in the REF, the basis for $\text{Nul } A$ will *not* consist of those columns—in fact, they will often be of the wrong size!

2 Find bases for $\text{Nul } A$ and $\text{Col } A$ for each matrix below:

(a) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ -1 & -1 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$

(d) $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ where $a \neq 0$

