

## Functions

### Collaborators:

As you transition to higher mathematics, you will see that math is often the study of functions. It is perhaps time, then, to define the word. Let  $S$  and  $T$  be sets. A **function** (or a **map**)  $f: S \rightarrow T$  is a rule that assigns to each element  $s \in S$  exactly one element  $f(s) \in T$ .

The element  $f(s) \in T$  is called the **image** of  $s$  under  $f$ . It is common say “ $f$  sends  $s$  to  $f(s)$ ” or “ $f$  maps  $s$  to  $f(s)$ ” and to write  $s \mapsto f(s)$ .

The set  $S$  is called the **domain** of  $f$ , and the set  $T$  is called the **codomain** of  $f$ . The set  $f(S) = \{f(s) \mid s \in S\}$ , the set of all possible values of  $f(s)$  with  $s \in S$ , is called the **image** (or the **range**) of  $f$ .

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $f(x) = x^2$ . Identify the domain, codomain, and image of  $f$ .

Notice that, in general, the range of  $f$  is a subset of the codomain of  $f$ . For some functions, the range will actually be the entire codomain. A function  $f: S \rightarrow T$  is called **onto** (or **surjective**) if  $f(S) = T$ . That is,  $f$  is onto if for all  $t \in T$ , there exists  $s \in S$  such that  $f(s) = t$ .

A function  $f: S \rightarrow T$  is called **one-to-one** (or 1-1 or **injective**) if no two elements of  $S$  are mapped to the same element of  $T$ . That is,  $f$  is one-to-one if  $f(s_1) = f(s_2)$  implies  $s_1 = s_2$ .

We say that  $f$  is **bijective** (or a bijection) if  $f$  is both onto and one-to-one.

**Examples.**

- (a) Consider the function  $f: \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  defined by  $f(m, n, r) = 2^m 3^n 6^r$ . Is  $f$  onto? Is  $f$  1-1?
- (b) Find an onto function from  $\mathbb{N}$  to  $\mathbb{Z}$ .

Given a function  $f: S \rightarrow T$  and an element  $s \in S$ , we may map  $s$  to  $f(s)$ . Since  $f(s)$  is simply some element of  $T$ , it is perhaps natural to consider mapping it again to some other set. If we have a function  $g: T \rightarrow U$ , we can apply  $f$  and then  $g$  to map  $s \mapsto f(s) \mapsto g(f(s))$ . The **composition** of  $f$  and  $g$  is the function  $g \circ f: S \rightarrow U$  defined by

$$(g \circ f)(s) = g(f(s)).$$

**Exercise.** Let  $S, T, U$  be sets and let  $f: S \rightarrow T, g: T \rightarrow U$  be functions. Though they can be found in the textbook, it is a good exercise to prove the following facts by yourself.

- (i) If  $f$  and  $g$  are both one-to-one, then so is  $g \circ f$ .
- (ii) If  $f$  and  $g$  are both onto, then so is  $g \circ f$ .
- (iii) If  $f$  and  $g$  are both bijective, then so is  $g \circ f$ .