



---

Representing Knot Groups Into  $SL(2, \mathbb{C})$

Author(s): D. Cooper and D. D. Long

Source: *Proceedings of the American Mathematical Society*, Vol. 116, No. 2 (Oct., 1992), pp. 547-549

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/2159765>

Accessed: 17-06-2016 21:40 UTC

## REFERENCES

Linked references are available on JSTOR for this article:

[http://www.jstor.org/stable/2159765?seq=1&cid=pdf-reference#references\\_tab\\_contents](http://www.jstor.org/stable/2159765?seq=1&cid=pdf-reference#references_tab_contents)

You may need to log in to JSTOR to access the linked references.

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<http://about.jstor.org/terms>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



*American Mathematical Society* is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*

## REPRESENTING KNOT GROUPS INTO $SL(2, \mathbb{C})$

D. COOPER AND D. D. LONG

(Communicated by Frederick R. Cohen)

**ABSTRACT.** We show that if a knot in  $S^3$  has nontrivial Alexander polynomial then the fundamental group of its complement has a representation into  $SL(2, \mathbb{C})$  whose image contains a free group of rank two.

Since the advent of Casson's invariant, one of the intriguing aspects of representations of the fundamental groups of three-dimensional  $\mathbb{Z}$ -homology spheres is the question of whether the group of every such homology sphere other than  $S^3$  has an irreducible representation into  $SU(2)$ . This is also of considerable relevance to Floer-Donaldson theory. The question in this generality seems to be some way from resolution (one corollary is the Poincaré conjecture!). However we may retreat to a weaker question by observing that if this were true then every homology sphere obtainable by surgery on a nontrivial knot in  $S^3$  has an irreducible representation into  $SL(2, \mathbb{C})$  and in particular, an affirmative answer implies that every nonabelian classical knot group has such a representation. This latter question also arises naturally in the context of [CS, CC].

In this note we prove the following

**Theorem 1.** *Let  $K$  be a knot in  $S^3$  whose Alexander polynomial is not identically 1. Then  $G(K) = \pi_1(S^3 \setminus K)$  has a representation into  $SL(2, \mathbb{C})$  whose image contains a free group of rank two.*

The reason for specifying that the image contain a free group rather than only requiring that it be nonsoluble is that we wish to rule out the rather uninteresting case of dihedral representations that although irreducible are guaranteed by the condition that  $\Delta_K(-1) \neq 1$ . Such representations are elliptic and already exist in  $SU(2)$ .

We also remark that the condition on the Alexander polynomial guarantees that  $G(K)$  has a representation into  $SL(2, \mathbb{C})$  that is soluble and nonabelian, (see [BZ] or [R]) and a natural way to try and prove Theorem 1 would be to show that any such representation can be deformed to an irreducible one. However our proof does not proceed in this way and it remains an interesting open question whether a knot in  $S^3$  can have all its soluble representations being isolated.

---

Received by the editors February 28, 1991.

1991 *Mathematics Subject Classification.* Primary 57M25.

The second author was partially supported by grants from the Sloan Foundation and the NSF.

©1992 American Mathematical Society  
0002-9939/92 \$1.00 + \$.25 per page

One interpretation of Theorem 1 is that the Alexander polynomial is a strictly weaker invariant of knottedness than the  $A$ -polynomial of [CC].

**Corollary 2.** *If  $K$  is a knot in  $S^3$  with nontrivial Alexander polynomial, then it has nontrivial  $A$ -polynomial.*

Various special cases are also of interest, for example we have

**Corollary 3.** *The group of every fibered knot in  $S^3$  has a representation into  $SL(2, \mathbb{C})$  whose image contains a free group of rank two.*

Corollary 2 follows immediately from Theorem 1, since the latter guarantees a representation that by [T], has to lie on a component of representations of complex dimension at least 1, so that there is at least one factor in the  $A$ -polynomial of  $K$ .

*Proof of Theorem 1.* We note that a calculation from the presentation (see [BZ]) reveals that the theorem is true for torus knots and it is certainly true for hyperbolic knots.

It follows from Thurston's hyperbolisation theorem that we can suppose  $K$  contains a nonboundary parallel incompressible embedded torus, so that  $K$  is a satellite knot. By Alexander's theorem, every torus embedded into  $S^3$  bounds a solid torus on at least one side and it follows that  $K$  contains an incompressible torus  $T$  that separates the knot complement into two pieces, one of which is an atoriodal knot complement in  $S^3$  and the other of which is a solid torus containing the knot  $K$ . Observe that since the exterior knot is atoriodal, it has a representation of the required sort.

As usual we define the algebraic (resp. geometric) winding numbers of  $K$  inside the solid torus to be the minimal number of algebraic (resp. geometric) intersections of  $K$  with a meridian disc of the solid torus. Note that the geometric winding is strictly larger than zero since  $T$  is incompressible.

There are now two cases. Suppose that the algebraic winding number of  $K$  is nonzero. Then we see easily that we may extend a random irreducible representation of the exterior knot across to  $K$  by using an abelian representation of  $\pi_1(D^2 \times S^1 \setminus K)$ ; the representation of  $G(K)$  so obtained satisfies the conclusions of the theorem provided the representation of the exterior knot did.

We are therefore reduced to the case that the algebraic winding number of  $K$  inside the solid torus is zero. Mark  $T$  using the longitude-meridian pair coming from the exterior knot. We may re-embed into  $S^3$  so that it is unknotted and so that the longitude on  $T$  now bounds a disc in  $S^3$ . The embedding yields a new knot, which we denote  $K_1$ . Notice that the winding number condition ensures that  $K$  has a Seifert surface lying entirely inside  $D^2 \times S^1$  and our choice of embedding implies that the Seifert form of  $K_1$  using this Seifert surface is the same as that of  $K$ . Whence  $K_1$  also has the same Alexander polynomial as  $K$ , and in particular  $K_1$  is knotted.

Further, there is a degree one map from the complement of  $K$  to the complement of  $K_1$  so that there is a surjective homomorphism  $G(K) \rightarrow G(K_1)$ , so that if we can prove the theorem for the knot  $K_1$ , we will be done.

We may repeat the above argument to obtain a chain of knots  $K, K_1, K_2, \dots$  all of which have the same Alexander polynomial, with the property that either  $K_n$  has a representation of the required sort and hence so does  $K$ , or we may form  $K_{n+1}$ . We conclude the proof then by a result of [S] where

it is shown that a chain of the above type cannot extend indefinitely without eventually terminating in the unknot. Since the Alexander polynomial condition excludes this last possibility, we must eventually reach a knot group for which we can find a representation and we are done.

## REFERENCES

- [BZ] G. Burde and H. Zeischang, *Knots*, de Gruyter Stud. Math., vol. 5, de Gruyter, Berlin, 1985.
- [CS] M. Culler and P. B. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. (2) **117** (1983), 109–146.
- [CC] D. Cooper, M. Culler, H. Gillet, D. D. Long, and P. B. Shalen, *Plane curves associated to character varieties of 3-manifolds*, preprint.
- [T] W. P. Thurston, *The geometry and topology of 3-manifolds*, preprint.
- [S] T. Soma, *On preimage knots in  $S^3$* , Proc. Amer. Math. Soc. **100** (1987), 589–592.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CALIFORNIA 93106