

# Nonabelian DT theory from abelian DT theory

Work in progress with Soheyla Feyzbakhsh

arXiv:2007.03037

arXiv:2103.02915

arXiv:21?? ??????

# Contents

- ▶ Joyce's generalised DT invariants
- ▶ The rough idea: how to pass from rank  $r$  to ranks  $\leq r - 1$
- ▶ Example: rank 1 to rank 0
- ▶ Results for rank  $\geq 1$
- ▶ Weak stability conditions
- ▶ Wall crossing
- ▶ Rank 0 to rank  $\pm 1$

## Preliminaries

Throughout we work on a fixed Calabi-Yau 3-fold  $X$   
(smooth complex projective variety with  $K_X \cong \mathcal{O}_X$ )  
with a fixed ample line bundle  $\mathcal{O}_X(1)$  and hyperplane class  
 $H := c_1(\mathcal{O}_X(1)) \in H^2(X, \mathbb{Z})$

satisfying the Bogomolov-Gieseker conjecture of Bayer-Macri-Toda  
(for which, see later) such as a quintic 3-fold (Chunyi Li).

Fix a Chern character  $c \in H^2(X, \mathbb{Q})$   
(or a numerical K-theory class  $c \in K_{\text{num}}(X)$ ).

Consider (semi)stable bundles, or sheaves, or complexes of sheaves  
 $E$  with  $\text{ch}(E) = c$ .

# Stability

There are many notions of stability for  $E$ .

The ones we consider can be written in terms of some central charge  $Z(\text{ch}(E)) \in \mathbb{C}$ .

Writing  $Z(E) = m(E) \exp(2\pi i\theta(E))$  we let the slope of  $E$  be  $\mu(E) := \tan \theta(E)$  and say  $E$  is (semi)stable if and only if

$$\mu(F) (\leq) \mu(E/F) \text{ for all nonzero } F \subsetneq E.$$

Here ( $\leq$ ) means  $<$  for stability and  $\leq$  for semistability. (Definition of  $F \subset E$  is tricky, but for now can just take subsheaves of sheaves.)

E.g.  $Z(E) = \int_X c_1(E) \cdot H^2 + i \text{rank}(E)$  gives  $\mu(E) = \frac{\text{deg}(E)}{\text{rank}(E)}$  and *slope stability*.

E.g.  $Z(E) = [\int_X \text{ch}(E(n)) \cdot \text{td}_X]_{\leq 2} + i \text{rank}(E)$  for large  $n \gg 0$  gives *Gieseker stability*.

## DT invariants

Choose  $c, H$  so that Gieseker semistability  $\implies$  Gieseker stability.  
(So all semistable sheaves have only scalar automorphisms.)

Then we can define an invariant  $DT(c) \in \mathbb{Z}$  “counting” Gieseker stable bundles or sheaves  $E$  with  $\text{ch}(E) = c$ .

(The moduli space  $M_c$  of Gieseker stable bundles is a projective scheme with “*perfect obstruction theory*” of virtual dimension zero.

Therefore it has a 0-dimensional virtual cycle, whose length is  $DT(c)$ .)

(Closely related to holomorphic bundles being the critical points of the holomorphic Chern-Simons functional.)

Can think of it as  $(-1)^{\dim M_c} e(M_c)$ .

Behrend showed each point  $E \in M_c$  can be assigned a multiplicity  $\chi^B(E) \in \mathbb{Z}$  such that  $DT(c)$  is the weighted Euler characteristic

$$e(M_c, \chi^B) = \sum_{i \in \mathbb{Z}} i e(\{\chi^B = i\}).$$

## Generalised DT invariants

For general  $c, H$  there are *strictly semistable* sheaves of charge  $c$ ; counting them is much more complicated.

Given stable objects of smaller charge, we can take all their direct sums (and extensions) to get semistable objects of charge  $c$  but large automorphism groups.

To invert this process Joyce and Kontsevich-Soibelman took a clever “*plethystic logarithm*” to get more controllable automorphism groups.

Joyce-Song were able to define a generalised invariant  $J(c) \in \mathbb{Q}$  which reduces to  $DT(c) \in \mathbb{Z}$  when semistable = stable.

Invariant under deformations of  $X$ .

Changes via a **wall-crossing formula** when we change the stability condition.

## The simplest wall crossing formula

Suppose a bundle  $F$  sits in an exact sequence

$$0 \longrightarrow A \longrightarrow F \longrightarrow B \longrightarrow 0 \quad (*)$$

with  $A, B$  stable, and that we can vary the stability condition so that the slopes of  $A$  and  $B$  cross.

Just below the wall ( $\mu(A) < \mu(B)$ )  $F$  will be stable.

Just above the wall  $F$  will be destabilised by  $(*)$ , but extensions in the opposite direction will become stable.

So on crossing the wall we lose a  $\mathbb{P}(\text{Ext}^1(B, A))$  of extensions  $(*)$  and gain a  $\mathbb{P}(\text{Ext}^1(A, B))$ .

So the Euler characteristic changes by  $-\text{ext}^1(B, A) + \text{ext}^1(A, B) = -\text{ext}^1(B, A) + \text{ext}^2(B, A) = \chi(B, A)$  by Serre duality. WCF is

$$J_+[F] = J_-[F] + (-1)^{\chi(B,A)-1} \chi(B, A) J[A] J[B].$$

## The rough idea

Fix  $n \gg 0$  so that  $H^{\geq 1}(E(n)) = 0$  for all semistable  $E$  of charge  $c$ .

Now replace  $E$  by the cokernel  $F$  of a section  $s \in H^0(E(n))$ ,

$$0 \longrightarrow \mathcal{O}(-n) \xrightarrow{s} E \longrightarrow F \longrightarrow 0.$$

Then  $\text{rank}(F) = \text{rank}(E) - 1$  and  $\text{ch}(F) = c_n := c - e^{-nH}$ .

To a **first approximation**, suppose all such  $E, F$  are stable for  $s \neq 0$ .

Then we find all  $F$ s come from an  $(E, s)$ , so  $M_{c_n}$  is a  $\mathbb{P}^{N-1}$ -bundle over  $M_c$  ( $N := \chi(E(n)) = \int_X c \cdot e^{nH} \cdot \text{td}_X$ ), so

$$J(c_n) = (-1)^{N-1} \cdot N \cdot J(c).$$

Now wall-cross to handle stability and get the correct formula....



## An example: rank 1 from rank 0

The rough idea actually works perfectly when  $\text{rank} = 1$ .

Here  $M_c$  is a moduli space of ideal sheaves  $E = \mathcal{I}_Z$ , where  $Z \subset X$  is a subscheme of dimension  $\leq 1$ . (Possibly tensored by a line bundle.)

Then  $s \in H^0(\mathcal{I}_Z(n)) \hookrightarrow H^0(\mathcal{O}(n))$  cuts out divisor  $\iota: D \hookrightarrow X$  and

$$F = \text{coker } s = \iota_*(\mathcal{I}_Z)$$

is a torsion sheaf supported on  $D$ . (“ $D4$ - $D2$ - $D0$  brane.”)

In this case  $E, F$  are Gieseker *stable* and are the only stable sheaves are of this form,

$$M_{c_n} \longrightarrow M_c \text{ is a } \mathbb{P}^{N-1}\text{-bundle, } N = \chi(c(n)),$$

and  $J(c_n) = (-1)^{N-1} \cdot N \cdot J(c)$ .

(Eg rank 2 bundles supported on  $D \in \left| \frac{n}{2}H \right|$  with  $\text{ch} = c_n$  are unstable.)

## GW invariants

By S-duality the D4-D2-D0 counts  $J(c_n)$  should have generating series which are vector-valued mock modular forms

([MSW97, dBCDMV06, GSY07, DM11, AMP19]; possibly need further wall-crossing to reach attractor stability)

whereas the abelian DT invariants  $J(c)$  count curves (and points) in  $X$  and — by the MNOP conjecture — can be written in terms of the Gromov-Witten invariants of  $X$ .

(Maulik-Nekrasov-Okounkov-Pandharipande conjecture now proved for most Calabi-Yau 3-folds by Pandharipande-Pixton.)

## Rank $r$ from rank 0

In higher rank  $r \geq 1$  there are corrections to the “*rough idea*”. They mean we can write rank  $r$  invariants in terms of rank  $r - 1, r - 2, \dots, 0$  invariants. Inductively we get to rank 0.

**Theorem** (arXiv:2103.02915)

For fixed  $c$  of rank  $\geq 1$ ,

$$J(c) = F(J(\alpha_1), J(\alpha_2), \dots)$$

is a universal polynomial in invariants  $J(\alpha_i)$ , with all  $\alpha_i$  of rank 0 and pure dimension 2.

So to express everything in terms of rank 1 (“*abelian*” theory) what’s left is to express **rank 0 in terms of rank 1**. (See later.)

(Formulae universal but not explicit. See recent “*Playing with the index of M-theory*” [DNPZ] for a prediction in noncompact toric case.)

## Weak stability conditions

We use the *weak stability conditions* of Bayer-Macri-Toda.

Pick  $b, w \in \mathbb{R}$  with  $w > \frac{1}{2}b^2$ .

Instead of  $\text{Coh}(X) \subset D(X)$  we work in the abelian category

$$\mathcal{A}_b := \left\{ E^{-1} \xrightarrow{d} E^0 : \mu_H^+(\ker d) \leq b, \mu_H^-(\text{coker } d) > b \right\}.$$

$\mu^+(F)$  is the maximum slope of a subsheaf of  $F$ ,

$\mu^-(F)$  is the minimum slope of a quotient sheaf of  $F$ .

On this we use the central charge

$$Z(E) = [\text{ch}_1(E) \cdot H^2 - b \text{ch}_0(E) H^3] + i[\text{ch}_2(E) \cdot H - w \text{ch}_0(E) H^3],$$

i.e. the slope function

$$\nu_{b,w}(F) = \begin{cases} \frac{\text{ch}_2(E) \cdot H - w \text{ch}_0(E) H^3}{\text{ch}_1(E) \cdot H^2 - b \text{ch}_0(E) H^3} & \text{if } \text{ch}_1(E) \cdot H^2 - b \text{ch}_0(E) H^3 \neq 0, \\ +\infty & \text{if } \text{ch}_1(E) \cdot H^2 - b \text{ch}_0(E) H^3 = 0. \end{cases}$$

## Bogomolov-Gieseker conjecture

We assume the *Bogomolov-Gieseker conjecture* of Bayer-Macri-Toda: a certain upper bound on  $\text{ch}_3$  for  $\nu_{b,w}$ -semistable objects  $E$ . Setting  $C_i := \text{ch}_i(E) \cdot H^{3-i}$ , it is

$$(C_1^2 - 2C_0C_2)w + (3C_0C_3 - C_1C_2)b + (2C_2^2 - 3C_1C_3) \geq 0,$$


It is a sufficient condition for the existence of *Bridgeland stability conditions* on  $X$ , and has now been proved for some Calabi-Yau 3-folds.

For instance Chunyi Li proved it for many  $(b, w)$  (enough for our applications) on quintic 3-folds  $X$ .


## Weak stability conditions II

Plot  $\Pi(E) := \left( \frac{\text{ch}_1(E) \cdot H^2}{\text{ch}_0(E) H^3}, \frac{\text{ch}_2(E) \cdot H^2}{\text{ch}_0(E) H^3} \right)$  on the same axes as  $(b, w)$ .

Then walls of instability for  $E$  become straight lines through  $\Pi(E)$  and  $\Pi(F)$ , where  $F$  is a destabilising sub- or quotient- object.



# Walls of instability for $c_n$



## Some aspects of the proof

- ▶ The Joyce-Song wall is where the  $\nu_{b,w}$ -slopes of  $F$  (of charge  $c_n$ ) and  $\mathcal{O}(-n)[1]$  coincide.
- ▶ Rotating the exact sequence  $0 \rightarrow \mathcal{O}(-n) \xrightarrow{s} E \rightarrow F \rightarrow 0$  in  $D(X)$  gives the destabilising exact triangle

$$E \rightarrow F \rightarrow \mathcal{O}(-n)[1].$$

- ▶ Below the wall  $F$  is destabilised by this, above the wall it is stable iff  $E$  is  $\nu_{b,w}$ -semistable and  $s$  does not factor through any semi-destabilising subsheaf.
- ▶ Gives wall-crossing formula
$$J_{b,w_+}(c_n) = J_{b,w_-}(c_n) + (-1)^{N-1} \cdot N \cdot J_{b,w}(c) + \dots$$
lower order terms from sections of destabilising subsheaves of  $E$  ( $\implies$  lower rank, so can induct on rank).
- ▶ Now wall cross **second term down to below the BG wall**, and all other terms **up to large volume chamber**.



## Some more aspects of the proof

- ▶ All these further wall crossings involve **only sheaves** – no more complexes of sheaves, nor shifts like  $\mathcal{O}(-n)[1]$ .
- ▶ These wall crossings spit out destabilising pieces which we also wall-cross up to the large volume chamber. Their wall-crossing also involves only sheaves. (So rank never increases.)
- ▶ At each stage the discriminant  $\Delta_H = (\text{ch}_1 \cdot H^2)^2 - 2(\text{ch}_2 \cdot H) \text{ch}_0 H^3$  decreases and cannot drop below 0.
- ▶ So a double induction on rank and  $\Delta_H$  turns  $J_{b,w_+}(c_n) = J_{b,w_-}(c_n) + (-1)^{N-1} \cdot N \cdot J_{b,w}(c) + \dots$  into  $J_{b,\infty}(c_n) = 0 + (-1)^{N-1} \cdot N \cdot J_{b,\infty}(c) + \dots$  with  $\dots$  of the form  $F(J_{b,\infty}(\alpha_i))$ ,  $\text{rank}(\alpha_i) \leq r - 1$
- ▶ A further wall-crossing passes from  $J_{b,\infty}$  to  $J$ .
- ▶ Thus have written  $J(c)$  in terms of  $J$  of lower rank sheaves.

## Ongoing work – rank 0

Now suppose  $c$  has rank 0.

Fix  $n \gg 0$  so that  $H^{\geq 1}(E(n)) = 0$  for all semistable  $E$  of charge  $c$ .

For a section  $s \in H^0(E(n))$ , again replace  $E$  by the *complex of sheaves*  $F \in D(X)$

$$F := \{\mathcal{O}(-n) \xrightarrow{s} E\}.$$

Since  $s$  is neither injective nor surjective  $F$  is no longer quasi-isomorphic to a sheaf (unlike when  $\text{rank}(E) > 0$ ).

Then  $\text{rank}(F) = -1$  and  $\text{ch}(F) = c_n := c - e^{-nH}$ .

## Rank 0 II

The shift by [1] of the *derived dual* of  $F$

$$F^\vee[1] := \{E^\vee \xrightarrow{s} \mathcal{O}(n)\}$$

has rank 1, and after wall crossing becomes a **stable pair**. After a further, older wall-crossing (Bridgeland, Toda) it becomes an **ideal sheaf**, recovering the MNOP (or GW) invariants again.

So the “*rough idea*” in this case gives a simple universal formula relating D4-D2-D0 counts to rank 1 DT invariants (curve counts), just as we wanted.

We haven't handled all of the corrections terms yet, but we have shown that when they involve other D4-D2-D0 branes, the D4 brane is a **sheaf** with **smaller** degree (intersection with  $H^2$ ). So we can set up an induction again.