

Name: Solution

Math 54, Spring 2009, Section 109  
Quiz 1

(1) Find the general solution:

$$\begin{array}{rcl} x & - 4y & + 2z = 3 \\ & 2y & + 6z = -8 \\ -2x & + 10y & + 2z = -14 \end{array}$$

$$\left[ \begin{array}{cccc} 1 & -4 & 2 & 3 \\ 0 & 2 & 6 & -8 \\ -2 & 10 & 2 & -14 \end{array} \right] \xrightarrow{\substack{R_1+R_3 \\ R_3}} \left[ \begin{array}{cccc} 1 & -4 & 2 & 3 \\ 0 & 2 & 6 & -8 \\ 0 & 2 & 6 & -9 \end{array} \right] \xrightarrow{-R_2+R_3 \rightarrow R_3} \left[ \begin{array}{cccc} 1 & -4 & 2 & 3 \\ 0 & 2 & 6 & -8 \\ 0 & 0 & 0 & -1 \end{array} \right]$$

$$\left[ \begin{array}{cccc} 1 & -4 & 2 & 3 \\ 0 & 2 & 6 & -8 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{\frac{1}{2}R_2 \rightarrow R_2} \left[ \begin{array}{cccc} 1 & -4 & 2 & 3 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{4R_2 + R_1 \rightarrow R_1} \left[ \begin{array}{cccc} 1 & 0 & 14 & -13 \\ 0 & 1 & 3 & -4 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$x_3$  free

$$x_1 = -13 - 14x_3$$

$$x_2 = -4 - 3x_3$$

or

$$\begin{bmatrix} -13 \\ -4 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -14 \\ -3 \\ 1 \end{bmatrix}$$

(2) (a) State the definition of a set  $\{\vec{v}_1, \dots, \vec{v}_p\}$  being linearly dependent.

The set is called linearly dependent if there exist scalars  $x_1, \dots, x_p \in \mathbb{R}$  such that  $x_1\vec{v}_1 + \dots + x_p\vec{v}_p = \vec{0}$ , and not all the  $x_i$  are 0.

(b) Can you have a set of two linearly dependent vectors in  $\mathbb{R}^4$ ? Give an example, or say why it is not possible.

Xer.

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 2 \\ 2 \end{bmatrix} \right\}$$

(3) For which values of  $h$  does the following vector equation have one solution? No solutions? Many solutions?

$$x_1 \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \\ -2 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 3 \\ -4 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ h \end{bmatrix}$$

Equivalent to  $\begin{bmatrix} 1 & 0 & 2 & 1 \\ 3 & -1 & 3 & 2 \\ 1 & -2 & -4 & h \end{bmatrix}$ 's system of equations.

$$\begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & -1 & -3 & -1 \\ 0 & -2 & -6 & h-1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 3 & 1 \\ 0 & -2 & -6 & h-1 \end{bmatrix} \rightarrow$$

$$\rightarrow \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & h+1 \end{bmatrix}$$

If  $h = -1$ , the system is consistent.

Since there is a free variable ( $x_3$ ), in

this case there are infinitely many solutions. If  $h \neq -1$ , then  
are no solutions. There is never just one solution.