

# MATH 164: HOMEWORK 7

Due Friday, May 22nd

Questions followed by \* are to be turned in. Questions without \* are extra practice. At least one extra practice question will appear on each exam.

## Question 1 (Textbook Problem 6.2.8)

---

Solve the following linear program using duality theory.

$$\begin{aligned} & \text{minimize } z = x_1 + 2x_2 + \cdots + nx_n, \\ & \text{subject to } x_1 \geq 1, \\ & \quad x_1 + x_2 \geq 2, \\ & \quad \cdots \\ & \quad x_1 + x_2 + \cdots + x_n \geq n, \\ & \quad x_1, x_2, \dots, x_n \geq 0. \end{aligned}$$

(Hint: show that  $x_1 = n$  and  $x_2, \dots, x_n = 0$  is feasible for the primal and find a feasible solution for the dual for which the values of the objective functions are equal. Then explain why this ensures you have found the optimal solution.)

## Question 2\* (Textbook Problem 6.2.16)

---

Consider the linear program

$$\begin{aligned} & \text{minimize } z = 2x_1 + 9x_2 + 3x_3, \\ & \text{subject to } -3x_1 + 2x_2 + x_3 \geq 1, \\ & \quad x_1 + 4x_2 - x_3 \geq 1, \\ & \quad x_1, x_2, x_3 \geq 0. \end{aligned}$$

- (i) Find the dual to this problem and solve it graphically.
- (ii) Use complementary slackness to obtain the solution to the primal.

## Question 3\*

---

Recall that a symmetric  $n \times n$  matrix  $A$  is **positive semi-definite** if  $x^T Ax \geq 0$  for all  $x \in \mathbb{R}^n$ . Show that a matrix  $A$  is positive semi-definite if and only if all eigenvalues of  $A$  are nonnegative. (Hint: Use the Spectral Theorem from linear algebra. This theorem tells you that if a matrix is symmetric, then there exists an orthonormal basis of eigenvectors.)

## Question 4\* (Similar to Textbook Problem 2.3.20)

---

Determine if the following functions are convex, concave, both, or neither.

- (a)  $f(x_1, x_2) = 2x_1 - 4x_2$
- (b)  $f(x_1, x_2) = x_1^2 + x_2^2$
- (c)  $f(x_1, x_2) = 2x_1^2 - 2x_1x_2 + 5x_2^2 + 31x_1 - 70x_2$

**Question 5\* (Similar to Textbook Problem 2.6.4)**

---

Consider the function

$$f(x_1, x_2) = 2x_1^3 + 2x_1x_2^2 + x_1 + 2x_2^3.$$

- (a) Find the first three terms of the Taylor series for  $f$  centered at  $x_0 = (2, 1)$ .
- (b) Evaluate this Taylor series for  $p = (-0.1, 0.1)^T$  and compare with the value of  $f(x_0 + p)$ . Would we expect these values to become more similar or more different if we took  $p = (-1, 1)^T$  instead?

**Question 6 (Textbook Problem 2.6.4)**

---

Find the first three terms of the Taylor series for

$$f(x_1, x_2) = \sqrt{x_1^2 + x_2^2}$$

about the point  $x_0 = (3, 4)^T$ .

**Question 7\* (Textbook Problem 2.6.6)**

---

Prove that if  $p^T \nabla f(x_k) < 0$ , then  $f(x_k + \epsilon p) < f(x_k)$  for  $\epsilon > 0$  sufficiently small. (Hint: Expand  $f(x_k + \epsilon p)$  as a Taylor series about the point  $x_k$  and look at  $f(x_k + \epsilon p) - f(x_k)$ . Pretend that  $p^T D^2(\xi)p$  is a constant independent of  $\epsilon$  and use the fact from class that  $C_1 \epsilon^2 < C_2 \epsilon$  for  $\epsilon$  sufficiently small.)

(While is not technically true that  $p^T D^2(\xi)p$  is a constant independent of  $\epsilon$ , those of you who took Math 131A could probably show that  $p^T D^2(\xi)p$  is bounded above by a constant  $M$ , which is all you actually need for this problem.)

**Question 8\* (Textbook Problem 3.2.2)**

---

Suppose  $A$  is an  $m \times n$  matrix with full row rank. We say that a matrix  $Z$  of dimension  $n \times r$ ,  $r \geq n - m$ , and rank  $n - m$  is a *null-space matrix* for  $A$  if it satisfies  $AZ = 0$ . If  $r = n - m$  (i.e. the columns of  $Z$  are linearly independent), then  $Z$  is a *basis matrix* for the null space of  $A$ .

Let  $Z$  be an  $n \times r$  null-space matrix for the matrix  $A$ . If  $Y$  is any invertible  $r \times r$  matrix, prove that  $\hat{Z} = ZY$  is also a null-space matrix for  $A$ . Clearly explain how you use that  $Y$  is invertible.