

## Negligible Subsets in the Space Of Homeomorphisms

Raymond Y Wong

We show that for various compact metric spaces  $X$ , the space of homeomorphisms  $H(X)$  is homeomorphic to  $H(X) \setminus K$ , where  $K = \bigcup_{i > 0} K_i \subset H(X)$  with each  $K_i$  is either (1) closed and equi-uniformly continuous or (2) topologically complete.

Our motivation for the study of negligible subsets is that, for a compact piecewise linear  $n$ -manifold  $M$ , it has been a long standing problem as to whether the space of homeomorphisms  $H(M)$  is an absolute neighborhood retract (The Homeomorphism Group Problem). It turns out ([GH]) that there is a dense  $G$  subset  $G \subset H(M)$  which is homeomorphic to a  $s$ -manifold, where  $s$  is the countable infinite product of open interval  $(-1, 1)$ , and such that the complement  $H(M) \setminus G$  is a countable union of closed sets  $\{K_i\}$  each of which is a  $Z$ -set in the sense of [An] (it means that, for any homotopically trivial open set  $U$  in  $H(M)$ ,  $U \setminus K_i$  remains homotopically trivial). It is therefore natural to ask whether the union  $\bigcup_{i > 0} K_i$  may be deleted from  $H(M)$ . If the answer is yes, then  $H(M)$  is homeomorphic to  $G$ .

**Notation.** For a compact metric space  $(X, d)$ , let  $C(X)$  denote the space of continuous functions of  $X$  into  $X$ . The metric defined on  $C(X)$  is  $(f, g) = \sup\{d(f(x), g(x)) \mid x \in X\}$ . Without loss of generality, we may assume  $(f, g) \leq 1$ . Let  $H(X)$  denote the subspace consisting of homeomorphisms of  $X$  onto  $X$ .  $H(X)$  is a topological group via compositions. Denote  $H^*(X) = \text{the closure of } H(X) \text{ in } C(X)$ . To say  $K_i \subset H(X)$  is

2000 Mathematics Subject Classification. 58D15, 57N20

Key words and phrases. Homeomorphisms, Space of homeomorphisms, Negligible subsets

equi-uniformly continuous we mean that, for any given  $\epsilon > 0$ , there exists a  $\delta_i > 0$ , such that for all  $f \in K$ ,  $d(f(x), f(y)) < \epsilon$  whenever  $d(x, y) < \delta_i$ . Finally  $K \subset H(X)$  is said to be negligible if  $H(X)$  is homeomorphic to  $H(X) \setminus K$ .

**Theorem 1.** Let  $X \subset \mathbb{R}^m$  be the  $m$  unit-ball of the Euclidean  $m$ -space  $\mathbb{R}^m$ . Denote by  $H(X)$  the subspace consisting of all  $f \in H(X)$  which is the identity on the boundary  $\partial X$  of  $X$ . Suppose  $K = \bigcup_{i>0} K_i \subset H(X)$ , where each  $K_i$  is closed and equi-uniformly continuous, then  $H(X)$  is homeomorphic to  $H(X) \setminus K$ .

**Proof.** The proof require the concept of Morse's  $\mu$ -length of paths ([Mo]): Let  $f : [a, b] \rightarrow X$  be a path in a metric space  $(X, d)$  where  $a < b$ . For each positive integer  $n$ , let  $A_n = \{(t_0, t_1, \dots, t_n) \mid a \leq t_0 \leq t_1 \leq \dots \leq t_n \leq b\}$ . For each  $(t_0, t_1, \dots, t_n) \in A_n$ , define  $\mu_n(f; t_0, t_1, \dots, t_n) = \max \{d(f(t_i), f(t_{i-1})) \mid i = 1, 2, \dots, n\}$  and

$$\mu_n(f) = \sup \{ \mu_n(f; t_0, t_1, \dots, t_n) \mid (t_0, t_1, \dots, t_n) \in A_n \}.$$

The  $\mu$ -length of  $f$  is 
$$\mu(f) = \lim_{n \rightarrow \infty} (1/2^n) \mu_n(f).$$

We first consider a single closed and equi-uniformly continuous subset  $K \subset H(X)$ .

Let  $J = [a, b] \subset X$  denote an interval with  $a < b$  and with  $a, b \in X$ . For each  $f \in K$ , by Lemma 1.3 of [Sa-Wo], there is a point  $t_f$ ,  $a < t_f < b$ , depending continuously on  $f$ , such that  $\mu(f|_{[a, t_f]}) = \mu(f|_{[t_f, b]})$ .

For any subset  $A \subset X$ , Denote  $\text{Mesh}(A) = \sup \{d(x, y) \mid x, y \in A\}$ . The main goal of the following lemma is to establish (d).

**Lemma 1.** (a) For any  $a < r < b$ ,  $\mu(f|_{[a, r]}) \leq \text{Mesh}(f([a, r]))$ .

(b) For any  $a < r < r' < b$ ,  $\mu(f|_{[a, r]}) < \mu(f|_{[a, r']})$ .

(c) Let  $E$  denote the identity map on  $X$ . then

$$\mu(E|_{[a, b]}) \leq \mu(f|_{[a, b]}) \leq 2\mu(f|_{[a, t_f]})$$

(d) there is a point  $r$ ,  $a < r < b$ , such that  $t_f > r$  for all  $f \in K$ .

**Proof.** (a) For any  $n$ ,  $(f|_{[a, r]}; t_0, t_1, \dots, t_n) \leq \text{Mesh}(f([a, r]))$ . Hence  $\mu_n(f) \leq \text{Mesh}(f([a, r]))$ . It follows that  $\mu(f|_{[a, r]}) \leq \text{Mesh}(f([a, r]))$ .

The first inequality of (c) is true since  $a, b \in X$  and each  $f \in K$  fixes the endpoints  $a, b$ . The rest of the proof (b) and (c) is a straightforward application of the definition of the  $\mu$ -length of  $f$  and the triangle inequality of the metric  $d$ .

(d) Let  $\epsilon = (1/3)\mu(E|_{[a, b]})$ . Since  $K$  is equi-uniformly continuous, there exists  $\delta > 0$ , such that for all  $f \in K$ ,  $d(f(x), f(y)) < \epsilon$  whenever  $d(x, y) < \delta$ . Consider a point  $r$ ,  $a < r < b$ , such that  $d(a, r) < \delta$ . Since the metric  $d$  is Euclidean,  $d(x, y) \leq d(a, r) < \delta$  for all  $x, y \in [a, b]$ . It follows for all  $f \in K$ ,  $d(f(x), f(y)) < \epsilon$ . Thus  $\mu(f|_{[a, r]}) \leq \text{Mesh}(f([a, r])) \leq \epsilon$  by Lemma 1(a). Using Lemma 1(c), we have  $\mu(f|_{[a, r]}) < \mu(f|_{[a, t_f]})$ . Hence  $r < t_f$ .

Inside the  $m$ -ball  $X$  we can construct a countable, mutually disjoint collection of intervals  $[a_i, b_i]$  (actually a line segment being regarded as an interval), with  $a_i, b_i \in X$  and the diameter  $(b_i - a_i)$  converges to 0. Mimicking the construction of  $f \in K$  on  $[a, b]$ , we consider the construction of  $f \in K$  on each  $[a_i, b_i]$ . Using lemma 1(d), we can choose, for each fix  $i$ , a point  $r_i$ ,  $a_i < r_i < b_i$ , such that  $t_{i_f} > r_i$  for all  $f \in K$ . Recall that  $t_{i_f}$  is the point in the interval  $(a_i, b_i)$  for which  $\mu(f|_{[a_i, t_{i_f}]}) = \mu(f|_{[t_{i_f}, b_i]})$ . Denote  $s = (a_i, b_i)$ .

Let  $H_0(X) = \{f \in H(X) \mid \text{for all } i, \mu(f|_{[a_i, m_i]}) = \mu(f|_{[m_i, b_i]})\}$ , where  $m_i = \text{mid-point of } [a_i, b_i]$ . As demonstrated in [Sa-Wo], there is a homeomorphism  $\theta : H(X) \rightarrow H_0(X) \times s$  taking  $f \in H(X)$  to the point  $\theta(f) = (f', t_f)$ , where  $f' \in H_0(X)$  and  $t_f = (t_{i_f})$ .

Thus for  $f \in K$ ,  $\theta(f) = (f', t_f)$  with each  $t_{i_f} > r_i > a_i$ . In other words, the image  $\theta(K)$  is a closed set whose projection into each factor  $(a_i, b_i)$  is contained in  $[r_i, b_i]$ . By the techniques of infinite-dimensional (I-D) topology ([An]), there is homeomorphism  $\eta : (H_0(X) \times s) \setminus K \rightarrow H_0(X) \times s$  and the homeomorphism changes only the  $s$ -coordinates of each point.

Now suppose  $K = \bigcup_{i>0} K_i$ . The fact that we can delete the infinite sequence  $\{K_i\}$  from  $H_0(X) \times s$  is also a result of I-D topology techniques. Basically we write  $s = s_1 \times s_2 \times \dots$ , an infinite product with each  $s_i$  a copy of  $s$ . We then delete each  $K_i$  from  $H_0(X) \times s$  changing only the  $s_i$ -coordinates of each point. Collectively we can construct a homeomorphism taking  $(H_0(X) \times s) \setminus K$  onto  $H_0(X) \times s$ .

**Theorem 2.** Let  $X$  be a compact metric space containing a closed neighborhood  $N$  homeomorphic to some  $k$ -simplex. For  $K \subset H(X)$ ,  $H(X)$  is homeomorphic to  $H(X) \setminus K$  provided  $K$  is a countable union of topologically complete subsets  $\bigcup_{i>0} K_i$ .

**Remark.** Employing Bessaga's ([Be]) approach, in [Do], Dobrowolski shows that for a compact subset  $K \subset E$ ,  $K$  is negligible in  $E$  for a list of spaces  $E$ . Including in the list is  $E = H(X)$ , where  $X$  is a locally compact space containing a bicollared set. Using different approach, a similar result on  $H(X)$  was proved by Mason ([Ma]). A key step in our argument required each  $K_i$  to be topologically complete, a condition weaker than compactness but not as weak as being locally homotopically trivial, which is required to settle the Homeomorphism Group Problem.

**Proof.** The proof is the results of the following three steps.

(1) *Construction of a pinching map  $\lambda$ .* The main idea is to construct a sequence of paths each starting from a point (identity) in  $H(X)$  and ending with a point (a pinching map) in  $H^*(X) \setminus H(X)$ . We use these paths to push  $\{K_i\}$  out of  $H(X)$ . First of all, inside  $N$  we may assume it contains an interval  $J = [a, b]$  with  $0 < b - a < 1$ . We choose a point  $r$ ,  $a < r < b$ . Starting with identity, the idea is to shrink the interval  $[r, b]$  onto the point  $b$ . In other words, we construct a path  $\{\tau_t\}$  satisfying (i) each  $\tau_t$  is the identity outside  $N$ , (ii)  $\tau_0 = \text{identity}$ ,  $\tau_t([a, b]) = [a, b]$ ,  $\tau_t|_N$  is a canonical piecewise linear map that shrinks the interval  $[r, b]$  onto the interval  $[(1 - t)r + tb, b]$  and (iii) for  $t < 1$ , each  $\tau_t \in H(X)$  and  $\tau_1$  collapses the entire interval  $[r, b]$  onto  $b$ .

Next along  $J$  we construct a countable, mutually disjoint collection of  $k$ -simplices  $\{N_i\}$ , each a smaller version of  $N$ , with  $\text{diameter}(N_i)$  converges to 0. For each  $i$

$> 0$ , we choose an interval  $[a_i, b_i] \subset N_i$  with  $a_i < b_i$ . We also choose, for each  $i$ , a point  $r_i$ ,  $a_i < r_i < b_i$ . We then construct a sequence of paths  $\{ \gamma_{it} \}_i$ , such that for each  $i$ , the homotopy  $\{ \gamma_{it} \}$  take place in  $N_i$  completely analogous to the homotopy  $\{ \gamma_t \}$  in  $N$ . Define  $\gamma : [0, 1] \rightarrow H^*(X)$  as follow. For each point  $t = (t_i)_{i>0}$ ,  $\gamma(t)$  is the map which is the identity outside  $\bigcup_{i>0} N_i$  and such that for each  $i$ ,  $\gamma(t)|_{N_i} = \gamma_{it}|_{N_i}$ . Clearly  $\gamma$  satisfies the following

- Lemma 2.** (i) Denote  $Q = \gamma([0, 1])$ . Then  $Q \subset H^*(X)$  and  $\gamma([0, 1]) \subset H(X)$ ,
- (ii) for  $\gamma = [0, 1] \setminus [0, 1)$ ,  $\gamma \subset H^*(X) \setminus H(X)$ ,
- (iii) for all  $t = (t_i)_{i>0}, t' = (t'_i)_{i>0}$ ,  $d(\gamma(t), \gamma(t')) = \sup_{i>0} |t_i - t'_i|$ ,
- (v)  $\gamma$  is an imbedding and therefore the images  $Q$  is an absolute

retract.

(2) *Construction of a contractive map  $\varphi$ .* We say a map  $\gamma : C(X) \rightarrow C(X)$  is contractive if there is some number  $0 \leq r < 1$ , such that  $d(\gamma(f), \gamma(g)) \leq r d(f, g)$  for all  $f, g \in C(X)$ . Now let  $K = \bigcup_{i>0} K_i \subset H(X)$  be given such that each  $K_i$  is a complete subset of  $H(X)$ . Let  $r = b - a < 1$ . For any  $f \in H^*(X)$ , denote  $t_i = 1 - r d(f, K_i)$ . Thus  $0 \leq t_i \leq 1$  (recall that we assume the metric  $d$  is bounded by 1) and that  $t_i = 1$  if and only if  $f \in K_i$ . Let  $t = (t_i)_{i>0}$ . Define  $\gamma : H^*(X) \rightarrow H^*(X)$  by  $\gamma(f) = \gamma(t)$ .

- Lemma 3.** (i)  $\gamma(H^*(X)) \subset Q (= \gamma([0, 1]))$ ,
- (ii)  $\gamma(K) \subset H^*(X) \setminus H(X)$ ,
- (iii)  $\gamma(H(X) \setminus K) \subset H(X)$  and
- (iv)  $\gamma$  satisfies  $d(\gamma(f), \gamma(g)) \leq r d(f, g)$ , for all  $f, g \in H^*$ .

Proof. (i)-(iii) is clear. To verify (iv), denote  $t = (t_i)_{i>0}$  and  $t' = (t'_i)_{i>0}$ , where  $t_i = 1 - r d(f, K_i)$  and  $t'_i = 1 - r d(g, K_i)$ . By Lemma 1(iii),  $d(\gamma(f), \gamma(g)) = d(\gamma(t), \gamma(t')) = \sup_{i>0} |t_i - t'_i| = \sup_{i>0} r |d(f, K_i) - d(g, K_i)|$ . Since  $|d(f, K_i) - d(g, K_i)| \leq d(f, g)$  for all  $i$ ,  $d(\gamma(f), \gamma(g)) \leq r d(f, g)$ .

(3) *A homeomorphism of  $H(X) \setminus K$  onto  $H(X)$ .* Let  $\gamma : H^*(X) \rightarrow H^*(X)$  be defined as in (2) above. Define  $\varphi : H(X) \setminus K \rightarrow H(X)$  by  $\varphi(f) = \gamma^{-1} f$ . We will show that  $\varphi$  is a

homeomorphism onto  $H$ . The condition that each  $K_i$  to be topologically complete is a key requirement to show that  $\pi$  is surjective.

First of all, the map  $\pi$  is well-defined since for  $f \in K$ ,  $\pi(f) \in H(X)$  (Lemma 3(iii)), so  $\pi(f)^{-1}$  exists. Composition and inverse operations in  $H(X)$  are continuous, so  $\pi$  is continuous. Secondly, it is straightforward to verify that  $\pi$  is invariant under right multiplication; that is, for any  $h \in H(X)$ ,  $\pi(fh) = (\pi(f)h)$  for all  $f, g \in H(X)$ .

To show  $\pi$  is one-to-one, suppose  $\pi(f) = \pi(g)$  for  $f, g \in H(X) \setminus K$ . Then  $\pi(f)^{-1}\pi(f) = \pi(g)^{-1}\pi(g)$ , or  $\pi(g)^{-1}\pi(f) = \pi(gf^{-1})$ . Denote  $e = \text{identity}$ . We have  $\pi(\pi(g)^{-1}\pi(f)) = \pi(\pi(g)^{-1}\pi(f)^{-1}\pi(f)) = \pi(\pi(gf^{-1}), e) = \pi(gf^{-1}, e) = \pi(g, f)$ . Since  $\pi$  is a contractive map (Lemma 3(iv)),  $f = g$ .

To show  $\pi$  is onto, let  $g_0 \in H(X)$  be given. Consider the map  $\pi : H^*(X) \rightarrow H^*(X)$  defined by  $\pi(f) = (f)g_0$ . By Lemma 3(i), the images  $\pi(H^*(X)) = (H^*(X))g_0 \in Qg_0$ . It follows that the restriction  $\pi|_{Qg_0} : Qg_0 \rightarrow Qg_0$ . Since  $Qg_0$  is an absolute retract by Lemma 2(iv),  $\pi|_{Qg_0}$  have a fixed point, say  $f_0 \in Qg_0$  such that  $\pi(f_0) = f_0$ .

We want to assert  $f_0 \in ( )g_0$  by showing that any  $f \in ( )g_0$  is not a fixed point of  $\pi|_{Qg_0}$ . Given any  $\epsilon > 0$ . Since  $( )g_0 \in H^*(X) \setminus H(X)$  and  $K_i \in H(X)$  is topologically complete,  $K_i$  is a closed relative to  $H^*$ . Thus any  $f \in ( )g_0$  must have a positive distance from  $K_i$ . Let  $t_i = 1 - r(f, K_i)$  and denote  $t = (t_i)_{i > 0}$ . It follows  $t \in [0, 1)$ . By Lemma 2(i),  $\pi(f) = (f)g_0 = (t)g_0 \in H(X)$ . In other words,  $\pi(f) \in f$  and so  $f_0 \in H(X)$ .

Next we assert that  $f_0 \in K$ . For if  $f_0 \in K$ , then  $f_0 = \pi(f_0) = (f_0)g_0 \in ( )g_0$ , a contradiction. Thus  $f_0 \in H(X) \setminus K$  and  $f_0 = \pi(f_0) = (f_0)g_0$ , or  $(f_0) = (f_0)^{-1}f_0 = g_0$ .

The verification that  $\pi^{-1}$  is continuous is rather straightforward and will be omitted. Thus  $\pi : H(X) \setminus K \rightarrow H(X)$  is a homeomorphism and the proof of Theorem 2 is complete.

### References

[An] Topological properties of the Hilbert cube and the infinite product of open intervals, Trans. Amer. Math. Soc. **126**(1967), 200-206.

- [Be] C. Bessage, Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys **26**(1978), 117-119.
- [Do] T. Dobrowolski, Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys **26**(1978), 535-545.
- [GH] R. Geoghegan and W. Haver, On the space of piecewise linear homeomorphisms of a manifold, Proc. Amer. Math. Soc. **55** (1976), 145-151.
- [Ma] W. Mason, The space of all self-homeomorphisms of a 2-cell which fix the cell's boundary is an absolute retract, Tran. Amer. Math. Soc., **161**(1971), 185-206.
- [Mo] M. Morse, A special parameterization of curves, Bull. Amer. Math. Soc., **42**(1936), 915-922.
- [Sa-Wo] On the space of Lipschitz homeomorphisms of a compact polyhedron, Pacific J of Math, **139**(1989), 195-207.

E-mail: [wong@math.ucsb.edu](mailto:wong@math.ucsb.edu)

Math Department, University of California, Santa Barbara, CA 93106