

Integral metaplectic modular categories

Adam Deaton* and Paul Gustafson†

*Department of Mathematics, Texas A&M University,
College Station, TX 77843, USA*

*deaton@math.tamu.edu

†pgustafs@math.tamu.edu

Leslie Mavrakis

*Department of Mathematics, Seattle Pacific University,
Seattle, WA 98119-1997, USA
mavrakisl@spu.edu*

Eric C. Rowell

*Department of Mathematics, Texas A&M University,
College Station, TX 77843, USA
rowell@math.tamu.edu*

Sasha Poltoratski

*Department of Mathematics,
University of California Berkeley,
Berkeley, CA 94720, USA
sashapol@gmail.com*

Sydney Timmerman‡

*Department of Physics and Astronomy,
Johns Hopkins University,
Baltimore, MD 21218-2686, USA
stimmer2@jhu.edu*

Benjamin Warren

*Department of Mathematics and Statistics,
Swarthmore College,
Swarthmore, PA 19081, USA
benjamindaviswarren@yahoo.com*

Qing Zhang

*Department of Mathematics, Texas A&M University,
College Station, TX 77843, USA
zhangqing2513@tamu.edu*

Received 28 January 2019

Accepted 10 March 2020

Published 8 May 2020

‡Corresponding author.

ABSTRACT

A braided fusion category is said to have Property **F** if the associated braid group representations factor through a finite group. We verify integral metaplectic modular categories have property **F** by showing these categories are group-theoretical. For the special case of integral categories \mathcal{C} with the fusion rules of $\mathrm{SO}(8)_2$ we determine the finite group G for which $\mathrm{Rep}(D^\omega G)$ is braided equivalent to $\mathcal{Z}(\mathcal{C})$. In addition, we determine the associated classical link invariant, an evaluation of the 2-variable Kauffman polynomial at a point.

Keywords: Group-theoretical braided fusion category; property **F**; link invariant; braid group representation.

Mathematics Subject Classification 2020: 18D10, 20F36, 17B37

1. Introduction

Bosonic topological phases of matter can be modeled by unitary modular categories [16], which may be regarded as the programming language for topological quantum computation. Physically speaking, every simple object corresponds to an anyon type, and the braid group representations provide quantum gates. When the braid group representation associated with each anyon type has finite image, a universal topological quantum computer cannot be constructed from any anyon type in \mathcal{C} using braiding alone. It is an important problem to characterize which categorical models admit such universal braiding gates, and which do not. A category is said to have Property **F** if the braid group representations associated with every simple object factor over a finite group [14]. The Property **F** conjecture states that a category has Property **F** if and only if it is weakly integral [14], i.e. if and only if every simple object X has $\dim(X)^2 \in \mathbb{Z}$.

Unitary modular categories with the same fusion rules as $\mathrm{SO}(N)_2$ for integer $N > 1$, or metaplectic modular categories, form an important class of (weakly integral) modular categories. As noted in [1], the study of metaplectic modular categories first arose from a generalization of Majorana zero modes. We focus on the case of integral metaplectic modular categories—those for which the dimension of every simple object is an integer.

A category \mathcal{C} that is group-theoretical, in some sense, “comes from” a finite group; the Drinfeld center $\mathcal{Z}(\mathcal{C})$ is braided equivalent to the representation category of the twisted Drinfeld double of a finite group G , $\mathrm{Rep}(D^w G)$ [7]. Most importantly, group-theoretical categories have Property **F** [8].

Theorem 1. *All integral metaplectic modular categories \mathcal{C} are group-theoretical.*

In particular, our work verifies the Property **F** conjecture for integral metaplectic modular categories.

Although this may seem to be a negative result from the perspective of topological quantum computation, there are two very compelling reasons to study (weakly) integral modular categories: (1) the non-Abelian anyons that are on everyone’s minds these days as being physically realizable are the Majorana anyons, which

are weakly integral and (2) universality may often be recovered by including non-topological operations [5].

The group-theoreticity of integral metaplectic modular categories suggests a couple further directions of investigation. One direction is determining the finite group G for which $\mathcal{Z}(\mathcal{C}) = \text{Rep}(D^w G)$. Another direction of particular interest is determining the link invariant associated with each of these categories. The extended Property **F** conjecture states that a category has Property **F** if and only if the associated link invariants are classical [15]. Any topological quantum computation using anyons from these categories will approximate their associated link invariants evaluated on a braid closure. Thus, understanding the link invariant is equivalent to understanding how a topological quantum computer created using these anyon systems will behave.

We examine the specific case of modular categories with the same fusion rules as $\text{SO}(8)_2$. This is the smallest case of an integral metaplectic modular category for which the Drinfeld center cannot be equivalent to the representation category of the untwisted double of any finite group; i.e. there must be a nontrivial cocycle. Additionally, all simple objects are of dimension either one or two, creating extra symmetry and removing ambiguity regarding the associated link invariant.

Theorem 2. *The link invariant associated with a category with fusion rules of $\text{SO}(8)_2$ is*

$$\frac{1}{2}(-1)^{w(D)} \sum_{X \subset L} (-i)^{\text{linking number}(X, L-X)},$$

where L is a link, D is its diagram, $w(D)$ is the writhe of D , and the sum is over components $X \subset L$.

In Sec. 2, we prove the group-theoreticity of integral metaplectic modular categories. Section 3 will contain our examination of categories with fusion rules of $\text{SO}(8)_2$.

2. Group-Theoreticity

Throughout we will assume that the reader is familiar with basic categorical notions, and refer to [2, 7] for the standard definitions and details. Before we proceed to our results we need some standard notation. Let S denote the S matrix of a braided fusion category \mathcal{C} . We say an objects X, Y in a braided fusion category \mathcal{C} centralize each other if $S_{XY} = \dim(X)\dim(Y)$, in other words the double braiding between X and Y is trivial: $c_{Y,X}c_{X,Y} = \text{Id}_{X \otimes Y}$. Strictly speaking this is a theorem in [13] characterizing the latter condition in terms of the S -matrix. For any subcategory \mathcal{L} of a braided fusion category \mathcal{C} the centralizer of \mathcal{L} denoted by $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})$ is the subcategory consisting of objects $Y \in \mathcal{C}$ that centralize all objects $X \in \mathcal{L}$. Let $\mathcal{L} \subset \mathcal{C}$. Then, the adjoint subcategory \mathcal{L}_{ad} is the smallest fusion subcategory of

\mathcal{C} that contains $X \otimes X^*$ for each simple object $X \in \mathcal{L}$, where X^* is the dual of X .

To prove group-theoreticity, we use the following key characterization.

Proposition 2.1 ([6, Corollary 4.14]). *A modular category \mathcal{C} is group-theoretical if and only if it is integral and there is a symmetric subcategory \mathcal{L} such that $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})_{\text{ad}} \subset \mathcal{L}$.*

Now consider an integral N -metaplectic modular category. There are three cases to consider depending on the value of N . The relevant fusion rules for each of these cases are listed below (Note: fusion rules for \sqrt{N} and $\sqrt{\frac{N}{2}}$ dimensional objects are not included, but can be found here [1, 3]).

As stated in [11], the unitary modular category $\text{SO}(N)_2$ for odd $N > 1$ has two simple objects, X_1, X_2 of dimension \sqrt{N} , two simple objects $\mathbf{1}, Z$ of dimension 1, and $\frac{N-1}{2}$ objects $Y_i, i = 1, \dots, \frac{N-1}{2}$ of dimension 2 [1]. Observe that, in order for a metaplectic modular category to be integral, N must be a perfect square. Observe that, in order for a metaplectic modular category to be integral, N must be a perfect square. The fusion rules are [1]:

- (1) $Z \otimes Y_i \cong Y_i, Z^{\otimes 2} \cong \mathbf{1};$
- (2) $Y_i \otimes Y_j \cong Y_{\min\{i+j, N-i-j\}} \oplus Y_{|i-j|}, \text{ for } i \neq j;$
- (3) $Y_i^{\otimes 2} = \mathbf{1} \oplus Z \oplus Y_{\min\{2i, N-2i\}}.$

The unitary modular category $\text{SO}(N)_2$ where $N \equiv 2 \pmod{4}$ has rank $k+7$, where $2k = N$. The group of isomorphism classes of invertible objects is isomorphic to \mathbb{Z}_4 . We will denote by g a generator of this group, and will abuse notation referring to the invertible objects as g^j . There are $k-1$ self-dual simple objects, Y_i , of dimension 2. The remaining four simples, V_i , have dimension \sqrt{k} . The fusion rules re-written from [11] are:

- (1) $g \otimes Y_i \cong Y_{k-i}, g^2 \otimes Y_i \cong Y_i;$
- (2) $Y_i^{\otimes 2} \cong \mathbf{1} \oplus g^2 \oplus Y_{\min\{2i, 2k-2i\}};$
- (3) $Y_i \otimes Y_j \cong Y_{\min\{i+j, 2k-i-j\}} \oplus Y_{|i-j|}, \text{ when } i+j \neq k;$
- (4) $Y_i \otimes Y_j \cong g \oplus g^3 \oplus Y_{|i-j|} \text{ when } i+j = k.$

The modular category $\text{SO}(N)_2$ with $N \equiv 0 \pmod{4}$ where $2k = N$, has rank $k+7$ and dimension $4N$ [11]. The simple objects have dimension 1, 2, and \sqrt{k} . All simple objects are self-dual, and the Y_i have dimension 2, while V_i, W_i have dimension \sqrt{k} . For $k > 2$ the key fusion rules re-written from [11] and are as follows:

- (1) $g \otimes Y_i \cong f \otimes Y_i \cong Y_{k-i};$
- (2) $Y_i^{\otimes 2} \cong \mathbf{1} \oplus f \oplus g \oplus fg, \text{ when } i = \frac{k}{2};$
- (3) $Y_i^{\otimes 2} \cong \mathbf{1} \oplus fg \oplus Y_{\min\{2i, 2k-2i\}}, \text{ when } i \neq \frac{k}{2};$
- (4) $Y_i \otimes Y_j \cong Y_{\min\{i+j, 2k-i-j\}} \oplus Y_{|i-j|}, \text{ when } i+j \neq k;$
- (5) $Y_i \otimes Y_j \cong g \oplus f \oplus Y_{|i-j|}, \text{ when } i+j = k.$

Proposition 2.2. *Let \mathcal{C} be a braided tensor category. \mathcal{C} is symmetric if and only if it coincides with its symmetric center $\mathcal{Z}_{\mathcal{C}}(\mathcal{C})$ [13].*

Proposition 2.3. *Let \mathcal{C} be a modular category and $\mathcal{L} \subset \mathcal{C}$ be a semisimple tensor subcategory. Then $\dim(\mathcal{L}) \dim(\mathcal{Z}_{\mathcal{C}}(\mathcal{L})) = \dim(\mathcal{C})$ [13].*

Definition 2.4. The balancing equation $\theta_i \theta_j S_{ij} = \sum_{k \in Irr(\mathcal{C})} N_{i^* j}^k d_k \theta_k$.

Lemma 2.5. *Every integral metaplectic modular category has a symmetric subcategory \mathcal{L} .*

- (i) For N odd, \mathcal{L} is generated by $\mathbf{1}, Z$ and Y_{it} where $t = \sqrt{N}$ and $1 \leq i \leq \frac{t-1}{2}$.
- (ii) For $N \equiv 2 \pmod{4}$, \mathcal{L} is generated by $\mathbf{1}, g^2$ and Y_{2ln} where $l = \sqrt{k}$ and $1 \leq n \leq \frac{l-1}{2}$.
- (iii) For $N \equiv 0 \pmod{4}$, \mathcal{L} is generated by $\mathbf{1}, f, g, fg$ and Y_{2lm} where $1 \leq m \leq \frac{l-2}{2}$.

Proof. First, we examine the case of N odd in detail. We need to show that \mathcal{L} is symmetric, that is, by Proposition 2.5, $\mathcal{L} = \mathcal{Z}_{\mathcal{L}}(\mathcal{L})$. By Proposition 2.6, $\dim(\mathcal{Z}_{\mathcal{C}}(\mathcal{L})) = 2t$.

There is a faithful \mathbb{Z}_2 grading on \mathcal{C} :

$$\mathcal{C}_1 = \{\mathbf{1}, Z, Y_i\}, \mathcal{C}_Z = \{X_i\}.$$

We know that $\mathcal{C}_{pt} \subset \mathcal{L}$. Applying the fact that $\mathcal{Z}_{\mathcal{C}}(C_{pt}) = \mathcal{C}_1$ [9], we know that $\mathcal{Z}_{\mathcal{C}}(\mathcal{L}) \subset \mathcal{C}_1$.

The object Z is a boson [1, Lemma 3.3]. We perform the de-equivariantization of \mathcal{C} by $\langle Z \rangle \cong \text{Rep}(\mathbb{Z}_2)$ as the trivial component of the de-equivariantization \mathcal{D}_0 , the image of \mathcal{C}_1 , is isomorphic to \mathbb{Z}_N [1, Lemma 3.4]. The de-equivariantization functor maps $\mathbf{1}$ and Z to 0, and each Y_i to the simple objects $i \oplus -i \in \mathbb{Z}_N$. The subcategory tensor generated by Y_i maps to $\langle i \rangle \in \mathbb{Z}_N$.

As $\langle t \rangle$ is the only subgroup of order t , it is the image of the only subcategory of dimension $2t$, which must be $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})$. Now it is clear $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})$ is the subcategory tensor generated by Y_t . Because \mathcal{L} is equal to its centralizer in \mathcal{C} , it is symmetric.

The proof for the case of $N \equiv 2 \pmod{4}$ proceeds in the same fashion. By Proposition 2.6, $\dim(\mathcal{Z}_{\mathcal{C}}(\mathcal{L})) = 4\ell$. The faithful grading is by \mathbb{Z}_4 :

$$\begin{aligned} \mathcal{C}_1 &= \{\mathbf{1}, g^2, Y_i\} \quad \text{where } i \text{ is even,} \\ \mathcal{C}_g &= \{V_1, V_4\}, \\ \mathcal{C}_{g^2} &= \{g, g^3, Y_i\} \quad \text{where } i \text{ is odd,} \\ \mathcal{C}_{g^3} &= \{V_2, V_3\}. \end{aligned}$$

We know that g^2 is a boson [4, Lemma 3.3]. As $g^2 \in \mathcal{L}$, $\mathcal{Z}_{\mathcal{C}}(\mathcal{L}) \subset \mathcal{Z}_{\mathcal{C}}(\langle g^2 \rangle)$. A dimension counting argument tells us that $\mathcal{Z}_{\mathcal{C}}(\langle g^2 \rangle)$ must be $\mathcal{C}_1 \cup \mathcal{C}_{g^2}$. Forming the de-equivariantization by g^2 , the image of $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})$ is contained within \mathcal{D}_0 . Following the proof from [1, Lemma 3.3] $\mathcal{D}_0 \cong \mathbb{Z}_N$, so it only contains the image of one

subcategory of the correct size and $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})$ is the subcategory tensor generated by Y_i . Therefore, $\mathcal{Z}_{\mathcal{L}}(\mathcal{L}) = \mathcal{L}$ and \mathcal{L} is symmetric.

The proof for the case of $N \equiv 0 \pmod{4}$ proceeds in exact fashion as the proof for $N \equiv 2 \pmod{4}$, except in this case the faithful grading is by $\mathbb{Z}_2 \times \mathbb{Z}_2$,

$$\mathcal{C}_1 = \{\mathbf{1}, f, g, fg, Y_i\} \quad \text{where } i \text{ is even,}$$

$$\mathcal{C}_g = \{V_1, V_2\},$$

$$\mathcal{C}_f = \{W_1, W_2\},$$

$$\mathcal{C}_{fg} = \{Y_i\} \quad \text{where } i \text{ is odd,}$$

we de-equivariantize by the boson fg [3, Lemma 4.3], and apply the balancing equation and dimension counting to show that $Y_i \in \mathcal{Z}_{\mathcal{C}}(fg)$, the preimage of \mathcal{D}_0 , for all i . \square

Theorem 2.6. *Integral metaplectic modular categories are group-theoretical.*

Proof. We examine the symmetric subcategory \mathcal{L} we defined for each case. Using the fusion rules for each case and $\mathcal{Z}_{\mathcal{C}}(\mathcal{L})$, which we found in the proof of our previous result, it is clear that $(\mathcal{Z}_{\mathcal{C}}(\mathcal{L}))_{\text{ad}} \subset \mathcal{L}$. Therefore, in each case \mathcal{C} is group-theoretical. \square

3. Categories With Fusion Rules of $\text{SO}(8)_2$

Categories with the fusion rules of $\text{SO}(8)_2$ present an interesting special case. In these categories, objects V_1, V_2, W_1, W_2 of dimension $\sqrt{\frac{N}{2}}$ have the same dimension as objects Y_i , creating extra symmetry.

Definition 3.1. A category \mathcal{C} is group-theoretical if and only if its Drinfeld center $\mathcal{Z}(\mathcal{C})$ is equivalent to the representation category of the twisted double of some finite group G :


$$\mathcal{Z}(\mathcal{C}) \cong \text{Rep } D^{\omega}(G);$$

Having determined that categories with the fusion rules of $\text{SO}(8)_2$ are group-theoretical, it is natural to ask which finite group they “come from.” Applying the facts that $|G|^2 = \dim(\text{Rep } D^{\omega}(G)) = \dim \mathcal{Z}(\mathcal{C}) = (\dim \mathcal{C})^2 = 1024$ and $\text{rank}(\text{Rep}(D^{\omega}G)) = \text{rank}(\mathcal{Z}(\mathcal{C})) = (\text{rank } \mathcal{C})^2 = 256$, we search the database of modular data for twisted Drinfeld double of finite groups of order less than 47 created by Gruen [10] for pairs (ω, G) that fit these restrictions. We determined that categories with the fusion rules of $\text{SO}(8)_2$ come from $G = [32, 49]$, the inner holomorph of the dihedral group of order 8. Notably, there are 72 Morita-equivalence classes of categories for which $\text{Rep}(D^{\omega}G)$ has the correct dimension and rank, and for each of these ω is a nontrivial cocycle.

4. Link Invariants

Since every topological quantum computation is equivalent to evaluating the link invariant associated with the anyonic system on some braid, another natural inquiry is to determine the link invariant associated with categories with fusion rules of $\mathrm{SO}(8)_2$. The 2-variable Kauffman polynomial is known to be associated with the quantum group $U_{q,\mathrm{so}(n)}$. Examining the work of Tuba and Wenzl, we know the invariant associated with these categories is Wenzl's parameterization of the 2-variable Kauffman polynomial, evaluated at $q = e^{\frac{i\pi}{8}}$, $r = -q^{-1}$ [17]. The extended Property **F** conjecture predicts that the link invariant associated with a braided fusion category \mathcal{C} with Property **F** is classical; therefore, we expect this evaluation to produce a classical invariant.

In the following definitions, D is the diagram of a link. D_+ , D_- , D_0 , and D_∞ are identical diagrams except near a point where they are



Definition 4.1. The semi-oriented Kauffman polynomial is a function

$$K : \{\text{Oriented links in } S^3\} \rightarrow \mathbb{Z}[a^{\pm 1}, z^{\pm 1}]$$

where $\mathbb{Z}[a^{\pm 1}, z^{\pm 1}]$ is the Laurent polynomial over a and z . It is uniquely defined by $K(L) = a^{-w(D)} \tilde{K}(D)$ where

$$\tilde{K} : \{\text{Oriented diagrams of links}\} \rightarrow \mathbb{Z}[a^{\pm 1}, z^{\pm 1}]$$

is uniquely determined by:

- (I) $\tilde{K}(\bigcirc) = 1$;
- (II) \tilde{K} is invariant under regular isotopy;
- (III) $a^{-1} \tilde{K}(\overset{|}{\circlearrowleft}) = \tilde{K}(D) = a \tilde{K}(\overset{|}{\circlearrowright})$;
- (IV) $\tilde{K}(D_+) + \tilde{K}(D_-) = z(\tilde{K}(D_0) + \tilde{K}(D_\infty))$.

Definition 4.2. Wenzl's parameterization of the Kauffman polynomial is

$$W : \{\text{Oriented links in } S^3\} \rightarrow \mathbb{Z}[r^{\pm 1}, q^{\pm 1}],$$

where $\mathbb{Z}[q^{\pm 1}, r^{\pm 1}]$ is the Laurent polynomial over r and q . It is uniquely defined by $W(L) = r^{-w(D)} \tilde{W}(D)$ where

$$\tilde{W} : \{\text{Oriented diagrams of links}\} \rightarrow \mathbb{Z}[r^{\pm 1}, q^{\pm 1}]$$

is uniquely determined by:

- (i) $\tilde{W}(\bigcirc) = 1$;
- (ii) \tilde{W} is invariant under regular isotopy;

- (iii) $r\tilde{W}(\overset{\circ}{\smile}) = \tilde{W}(|) = r^{-1}\tilde{W}(\smile);$
(iv) $\tilde{W}(D_-) - \tilde{W}(D_+) = (q - q^{-1})(\tilde{W}(D_0) - \tilde{W}(D_\infty)).$

Definition 4.3. The Dubrovnik parameterization of the Kauffman polynomial is

$$F : \{\text{Oriented links in } S^3\} \rightarrow \mathbb{Z}[r^{\pm 1}, q^{\pm 1}],$$

where $\mathbb{Z}[q^{\pm 1}, r^{\pm 1}]$ is the Laurent polynomial over r and q . It is uniquely defined by $F(L) = r^{-w(D)} \tilde{F}(D)$ where

$$\tilde{F} : \{\text{Oriented diagrams of links}\} \rightarrow \mathbb{Z}[\alpha^{\pm 1}, \omega^{\pm 1}]$$

is uniquely determined by:

- (i) $\tilde{F}(\bigcirc) = 1$;
 - (ii) \tilde{F} is invariant under regular isotopy;
 - (iii) $\alpha^{-1}\tilde{F}(\overset{\wedge}{\bigcirc}) = \tilde{F}(|) = \alpha\tilde{F}(\overset{\wedge}{|})$;
 - (iv) $\tilde{F}(D_+) - \tilde{F}(D_-) = \omega(\tilde{F}(D_0) - \tilde{F}(D_\infty))$.

Theorem 4.4. The link invariant associated with categories with the same fusion rules as $\mathrm{SO}(8)_2$ is classical and given by

$$[W(L)]_{(r,q)=\left(-\frac{1}{q}, e^{\frac{i\pi}{8}}\right)} = \frac{1}{2} \sum_{X \subset L} i^{\text{linking number}(X, L-X)}.$$

Proof. As $N = 8$, we know that the link invariant associated with these categories is $W(L)$ with $r = q^7$ [17]. Lemma 7.5 of [17] shows that Birman–Murakami–Wenzl algebras $\text{BMW}(q, q)$, the algebras associated with the 2-variable Kauffman polynomial, do not depend on the particular value of q . Choosing $q = e^{\frac{i\pi}{8}}$, we have $r = -\frac{1}{q}$.

The only differences between the various parameterizations of the 2-variable Kauffman polynomial lie in their twist and skein relations. It is easy to see

$$W(L) = [F(L)]_{(\alpha,\omega)=(r^{-1},-(q-q^{-1}))}.$$

Applying [12, Theorem 1], then

$$\begin{aligned} [W(L)]_{(r,q)=\left(-\frac{1}{q}, e^{\frac{i\pi}{8}}\right)} &= (-1)^{c(L)-1} [K(L)]_{(a,z)=(ir^{-1}, i(q-q^{-1}))} \\ &= (-1)^{c(L)-1} [K(L)]_{(a,z)=(-q^5, q^5+(q^5)^{-1})}. \end{aligned}$$

Then, taking an expression for this evaluation of $K(L)$ of [12, Table 1] we have

$$[W(L)]_{(r,q)=\left(-\frac{1}{q}, e^{\frac{i\pi}{8}}\right)} = \frac{1}{2} \sum_{X \subset L} i^{\text{linking number}(X, L-X)}$$

which is quite classical indeed.

Acknowledgment

This research was carried out as part of an REU at Texas A&M University in which L.M., S.P., S.T., and B.W. participated. This research of E.C.R., A.D., P.G., and Q.Z. was partially supported by NSF grant DMS-1664359, and that of L.M., S.P., S.T., and B.W. by NSF grant DMS-1757872.

References

- [1] E. Ardonne, M. Cheng, E. C. Rowell and Z. Wang, Classification of metaplectic modular categories, *J. Algebra* **466** (2016) 141–146.
- [2] B. Bakalov and A. Kirillov Jr, *Lectures on Tensor Categories and Modular Functors*, University Lecture Series, Vol. 21 (American Mathematical Society, Providence, RI, 2001).
- [3] P. Bruillard, P. Gustafson, J. Y. Plavnik and E. C. Rowell, Dimension as a quantum statistic and the classification of metaplectic categories, in *Topological Phases of Matter and Quantum Computation*, Contemp. Math., Vol. 747 (Amer. Math. Soc., Providence, RI, 2020), pp. 89–113.
- [4] P. Bruillard, J. Y. Plavnik and E. C. Rowell, Modular categories of dimension p^3m with m square-free, *Proc. Amer. Math. Soc.* **147**(1) (2019) 21–34.
- [5] S. X. Cui, S.-M. Hong and Z. Wang, Universal quantum computation with weakly integral anyons, *Quantum Inf. Process.* **14**(8) (2015) 2687–2727.
- [6] V. Drinfeld, S. Gelaki, D. Nikshych and V. Ostriker, Group-theoretical properties of nilpotent modular categories, Preprint (2007), arXiv:0704.0195.
- [7] P. Etingof, D. Nikshych and V. Ostriker, On fusion categories, *Ann. Math.* **162** (2005) 581–642.
- [8] P. Etingof, E. C. Rowell and S. Witherspoon, Braid group representations from twisted quantum doubles of finite groups, *Pacific J. Math.* **234**(1) (2008) 33–41.
- [9] S. Gelaki and D. Nikshych, Nilpotent fusion categories, *Adv. Math.* **217**(3) (2008) 1053–1071.
- [10] A. Gruen, Computing modular data for pointed fusion categories, Preprint (2018), arXiv:1808.05060.
- [11] P. Gustafson, E. C. Rowell and Y. Ruan, Metaplectic categories, gauging, and property F, to appear in *Tohoku Math. J.*, Preprint (2018), arXiv:1808.00698.
- [12] W. B. R. Lickorish, Some link-polynomial relations, *Math. Proc. Cambridge Philos. Soc.* **105**(1) (1989) 103–107.
- [13] M. Müger, On the structure of modular categories, *Proc. London Math. Soc.* (3) **87**(2) (2003) 291–308.
- [14] D. Naidu and E. C. Rowell, A finiteness property for braided fusion categories, *Algebr. Represent. Theory* **14**(5) (2011) 837–855.
- [15] E. C. Rowell, Two paradigms for topological quantum computation, in *Advances in Quantum Computation*, Contemp. Math., Vol. 482 (American Mathematical Society, Providence, RI, 2009), pp. 165–177.
- [16] E. C. Rowell and Z. Wang, Mathematics of topological quantum computing, *Bull. Amer. Math. Soc.* **55** (2018) 183–238.
- [17] I. Tuba and H. Wenzl, On braided tensor categories of type BCD, *J. Reine Angew. Math.* **581** (2005) 31–69.
- [18] H. Wenzl, Quantum groups and subfactors of type B, C, and D, *Comm. Math. Phys.* **133**(2) (1990) 383–432.