

MATH 3B Final practice problems

1. A honeybee population starts with 100 bees and increases at a rate of $n'(t)$ bees per week. What does $100 + \int_0^4 v'(t) dt$ represent?

Solution: By FTC2/Net Change theorem,

$$100 + \int_0^4 v'(t) dt = 100 + n(4) - n(0) = 100 + n(4) - 100 = n(4),$$

and $n(4)$ is just the bee population after 4 weeks.

2. Compute the following:

(a) $\int e^{-x} \cos(2x) dx$

Solution: Apply integration by parts twice and get $\int e^{-x} \cos(2x) dx$ by itself.

(b) $\int \frac{1+x^2}{\sqrt{3x+x^3}} dx$

Solution: Let $u = 3x + x^3$. Then $\frac{1}{3} du = (1+x^2) dx$ and the rest is straightforward.

(c) $\int \cos(\sqrt{x}) dx$

Solution: Let $u = \sqrt{x}$. Then $du = \frac{1}{2\sqrt{x}} dx \Rightarrow 2\sqrt{x} du = 2udu = dx$. Then integration by parts.

(d) $\frac{d}{dx} \int_x^{x^2} t^3 dt$

Solution: By FTC1 and the chain rule,

$$\frac{d}{dx} \int_x^{x^2} t^3 dt = \frac{d}{dx} \left[\int_0^{x^2} t^3 dt - \int_0^x t^3 dt \right] = (x^2)^3 2x - x^3 = 2x^7 - x^3.$$

(e) $\int \frac{(\ln(x))^3}{x} dx$

Solution: Let $u = \ln(x)$, so $du = \frac{1}{x} dx \dots$

(f) $\int \frac{x^3 - \sqrt{x}}{x^2} dx$

Solution: Write as two fractions and integrate: $\frac{x^3 - \sqrt{x}}{x^2} = x - x^{-\frac{3}{2}}$.

(g) $\int \sin^3(x)\cos^2(x) dx$

Solution: $\int \sin^3(x)\cos^2(x) dx = \int (1 - \cos^2(x))\sin(x)\cos^2(x) dx$.
Let $u = \cos(x) \Rightarrow -du = \sin(x)dx$...

(h) $\int_0^1 \frac{4}{4x - 1} dx$

Solution: Notice that the integrand has a discontinuity at $x = 1/4$, so

$$\int_0^1 \frac{4}{4x - 1} dx = \lim_{t \rightarrow 1/4^-} \int_0^t \frac{4}{4x - 1} dx + \lim_{t \rightarrow 1/4^+} \int_t^1 \frac{4}{4x - 1} dx.$$

If you show that one of these integrals diverge, then the whole thing does (both diverge).

(i) $\int \frac{1}{x^2 - 1} dx$

Solution: Factor and do partial fractions...

3. Consider the region $R = \{(x, y) : x \geq 1, 0 \leq y \leq 1/x\}$.

(a) Show that the region R has infinite area.

Solution: The area of the region R is

$$\int_1^\infty 1/x dx = \lim_{t \rightarrow \infty} \int_1^t 1/x dx = \lim_{t \rightarrow \infty} [\ln(x)]_1^t = \lim_{t \rightarrow \infty} \ln(t) - 0 = \infty.$$

(b) Find the volume of the solid obtained by rotating R about the x -axis. It is finite, which is kinda crazy given part (a), huh?

Solution: Method 1: Disks

$$\pi \int_1^\infty r^2 dx = \pi \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2} dx = \pi \lim_{t \rightarrow \infty} \left[-\frac{1}{x} \right]_1^t = \pi(0 - (-1)) = \pi.$$

Solution: Method 2: Shells

$$2\pi \int_0^1 r h dy = 2\pi \int_0^1 y \left(\frac{1}{y} - 1 \right) dy = 2\pi \int_0^1 1 - y dy = 2\pi \left[y - \frac{y^2}{2} \right]_0^1 = \pi.$$

Notice that for the first method we had an improper integral, and for the second we didn't, but we did need to solve for x in terms of y .

4. Find the average value of $f(x) = \frac{x}{x-6}$ for $0 \leq x \leq 2$.

Solution 1: Making the substitution $u = x - 6 \Rightarrow x = u + 6, du = dx$ we have

$$\begin{aligned} f_{avg} &= \frac{1}{2-0} \int_0^2 \frac{x}{x-6} dx = \frac{1}{2} \int_{-6}^{-4} \frac{u+6}{u} du = \frac{1}{2} \int_{-6}^{-4} 1 + \frac{6}{u} du = \frac{1}{2} [u + 6\ln|u|]_{-6}^{-4} \\ &= \frac{1}{2} [x - 6 + 6\ln|x - 6|]_0^2 = \frac{1}{2} [(2 - 6 + 6\ln(4)) - (0 - 6 + 6\ln(6))] = \frac{1}{2} [2 + 6\ln(4) - 6\ln(6)]. \end{aligned}$$

Solution 2: Wouldn't it be nice if the numerator and denominator were the same?

$$\begin{aligned} f_{avg} &= \frac{1}{2-0} \int_0^2 \frac{x}{x-6} dx = \frac{1}{2} \int_0^2 \frac{x-6+6}{x-6} dx = \frac{1}{2} \int_0^2 1 dx + \frac{1}{2} \int_0^2 \frac{6}{x-6} dx = \frac{1}{2} [x + 6\ln|x - 6|]_0^2 \\ &= \frac{1}{2} [(2 + 6\ln(4)) - (0 + 6\ln(6))] = \frac{1}{2} [(2 + 6\ln(4)) - 6\ln(6)]. \end{aligned}$$

5. Find the work required to lift a chain lying on the ground to a height of 10 feet if the chain weighs 20 pounds and is 10 feet long.

Solution: Method 1: Chop up the object. We chop up the chain into pieces of length Δx ft. Each piece weighs Δx ft \times 2 lb/ft = $2\Delta x$ lbs. A piece x ft above ground needs to be lifted up another $10 - x$ ft, so the work required for each piece of chain is $2(10 - x)\Delta x$. Adding up the work required for each piece gives $\int_0^{10} 2(10 - x)dx = 100J$.

Solution: Method 2: Chop up the distance. Let x be the height above the ground. We'll add up the work required to lift the chain each tiny distance Δx . If we're at a height of x feet, we have to lift x feet of chain and the rest is coiled up on the ground. The force to move each tiny distance = x feet \times 2 lb/ft = $2x$ lb, so the total work is $W = \int_0^{10} 2x dx = 100J$.

6. Find the length of the arc $y = x^{3/2}$ for $0 \leq x \leq 4$.

Solution: $\frac{dy}{dx} = \frac{3}{2}x^{1/2}$, so $L = \int_0^4 \sqrt{1 + \frac{9}{4}x} dx$. Let $u = 1 + \frac{9}{4}x \Rightarrow \frac{4}{9}du = dx$; the rest should be simple enough.

7. Evaluate $\int_0^\infty \frac{1}{x-2} dx$ or show that the integral diverges.

Solution: Notice we are integrating over an infinite interval and $f(x) = \frac{x}{x-2}$ is discontinuous at $x = 2$, which is in our interval of integration. So we break up the integral as

$$\int_0^\infty \frac{1}{x-2} dx = \lim_{t \rightarrow 2^-} \int_0^t \frac{1}{x-2} dx + \lim_{t \rightarrow 2^+} \int_t^3 \frac{1}{x-2} dx + \lim_{t \rightarrow \infty} \int_3^t \frac{1}{x-2} dx.$$

If one of the integrals on the right diverges, then $\int_0^\infty \frac{1}{x-2} dx$ diverges. The last one looks a lot like $\int \frac{1}{x} dx$ so we show it diverges:

$$\lim_{t \rightarrow \infty} \int_3^t \frac{1}{x-2} dx = \lim_{t \rightarrow \infty} [\ln|x-2|]_3^t = \lim_{t \rightarrow \infty} \ln|t-3| - 0 = \infty.$$

Thus $\int_0^\infty \frac{1}{x-2} dx$ diverges.

8. Find the volume of the solid obtained by rotating the region bounded by the curves $y = 2x$ and $y = x^2$ about the x -axis.

Solution: The curves intersect at $x = 0$. We find the other intersection point: $2x = x^2 \Rightarrow x = 2$. Graph them or note that $2x \geq x^2$ on $[0, 2]$. Washers seem the best choice as then we'll integrate with respect to x and we already have y in terms of x :

$$\pi \int_0^2 r_o^2 - r_i^2 dx = \pi \int_0^2 (2x)^2 - (x^2)^2 dx = \pi \int_0^2 4x^2 - x^4 dx = \pi \left[\frac{4x^3}{3} - \frac{x^5}{5} \right]_0^2 = \pi \left(\frac{32}{3} - \frac{32}{5} \right).$$

9. Let $f(x) = x^2$. Approximate $\int_0^4 x^2 dx$ using a right hand Riemann sum and 4 subintervals.

Solution: We approximate the area by adding up the area of 4 rectangles, whose width is $\Delta x = \frac{b-a}{n} = \frac{4-0}{4} = 1$, and whose height is determined by the value of $f(x)$ at the right endpoints of each rectangle, $x = 1, 2, 3$, and 4 . This yields

$$\int_0^4 x^2 dx \approx \Delta x [f(1) + f(2) + f(3) + f(4)] = (1)[1^2 + 2^2 + 3^2 + 4^2] = 30.$$

10. Find the arc length function $s(x)$ from the point $(1, 0)$ for the function $y = \int_1^x \sqrt{t^3 - 1} dt$.

Solution: By FTC1, $\frac{dy}{dx} = \frac{d}{dx} \int_1^x \sqrt{t^3 - 1} dt = \sqrt{x^3 - 1}$, so

$$\sqrt{1 + [dy/dx]^2} = \sqrt{1 + x^3 - 1} = \sqrt{x^3} = x^{3/2}, \text{ and } s(x) = \int_1^x t^{3/2} dt = \frac{2}{5} [t^{5/2}]_1^x = \frac{2}{5} (x^{5/2} - 1).$$

11. Hydrostatic force and center of mass: you just did a homework on it :)